

# A new critical exponent $\varphi$ and its logarithmic counterpart $\hat{\varphi}$

R. Kenna<sup>1</sup>, B. Berche<sup>2</sup>

<sup>1</sup> Applied Mathematics Research Centre, Coventry University, Coventry, CV1 5FB, England

<sup>2</sup> Statistical Physics Group, Institut Jean Lamour, UMR CNRS 7198, Université de Lorraine, B.P. 70239, 54506 Vandœuvre lès Nancy Cedex, France

Received December 12, 2012, in final form March 11, 2013

It is well known that standard hyperscaling breaks down above the upper critical dimension  $d_c$ , where the critical exponents take on their Landau values. Here, we show that this is because in standard formulations in the thermodynamic limit, distance is measured on the correlation-length scale. However, the correlation-length scale and the underlying length scale of the system are not the same at or above the upper critical dimension. Above  $d_c$  they are related algebraically through a new critical exponent  $\varphi$ , while at  $d_c$  they differ through logarithmic corrections governed by an exponent  $\hat{\varphi}$ . Taking proper account of these different length scales allows one to extend hyperscaling to all dimensions.

**Key words:** hyperscaling, critical dimension, correlation length, critical exponents, scaling relations

**PACS:** 64.60.-i, 64.60.an, 05.50.+q, 64.60.De, 11.10.Kk

## 1. Introduction

Since the 1960's, the scaling relations between critical exponents have been of fundamental importance in the theory of critical phenomena [1–3]. Six primary critical exponents,  $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\delta$ ,  $\eta$  and  $\nu$ , have played the most important roles and these are related by four famous scaling relations. One of these — the hyperscaling relation — involves the dimensionality  $d$  of the system. It has long been known that hyperscaling, in its standard form, fails above the upper critical dimension  $d = d_c$  where the critical exponents take their Landau, mean-field values. E.g., for the Ising model above  $d_c = 4$ , one has  $\alpha = 0$ ,  $\beta = 1/2$ ,  $\gamma = 1$ ,  $\delta = 3$ ,  $\eta = 0$  and  $\nu = 1/2$ , irrespective of the dimensionality  $d$ . Here, we report on a more complete form for the hyperscaling relation which holds in all dimensions [4]. This involves a new critical exponent which we denote by  $\varphi$  (pronounced “koppa” [5]) and which characterises the finite-size scaling (FSS) of the correlation length. We report the evidence for the universality of  $\varphi$  through numerical studies of the five-dimensional Ising model with free boundary conditions [4].

We also examine the hyperscaling at the upper critical dimension, which is characterised by multiplicative logarithmic corrections. These corrections are also characterised by critical exponents which have scaling relations between them. The logarithmic hyperscaling relation involves an exponent  $\hat{\varphi}$  which is the logarithmic analogue of  $\varphi$  [6, 7].

We consider a lattice spin system in  $d$  dimensions. In units of the lattice constant, its linear extent is  $L$ . We denote by  $P_L(t)$  the value of a function  $P$  measured on such a system at a reduced temperature  $t$ . The latter is defined as

$$t = \frac{T - T_L}{T_L}, \quad (1.1)$$

where  $T_L$  is the value of the temperature  $T$  at which the finite-size reduced susceptibility (defined below) peaks and is referred to as the pseudocritical point.

In the infinite-volume limit,  $T_L$  becomes the critical point  $T_\infty \equiv T_c$  and the specific heat and correlation length scale nearby as

$$c_\infty(t) \sim t^{-\alpha}, \quad \xi_\infty(t) \sim t^{-\nu}. \quad (1.2)$$

The standard form of the hyperscaling relation, which is valid at and below the upper critical dimension, links the critical exponents in equation (1.2),

$$\nu d = 2 - \alpha. \quad (1.3)$$

Equation (1.3) was proposed by Widom [8, 9]. Kadanoff later presented an alternative but similar argument for it [10] and Josephson derived the related inequality  $\nu d \geq 2 - \alpha$  on the basis of plausible but non-rigorous assumptions [11, 12]. The scaling relations, including equation (1.3), are now well understood through the renormalization group [13].

The Landau or mean-field values  $\alpha = 0$  and  $\nu = 1/2$  for the Ising model are well established for all values of  $d$  at and above the upper critical dimension  $d_c = 4$ . Since  $\alpha$  and  $\nu$  are fixed for  $d > d_c$ , equation (1.3) cannot hold there. This is referred to as the collapse of hyperscaling in high dimensions.

Here, we introduce a new critical exponent  $\varphi$  which characterises the leading FSS of the correlation length,

$$\xi_L(0) \sim L^\varphi. \quad (1.4)$$

We show that the incorporation of  $\varphi$  into the hyperscaling relation (1.3) via

$$\frac{\nu d}{\varphi} = 2 - \alpha, \quad (1.5)$$

renders it valid in all dimensions (with  $\varphi = 1$  in  $d \leq d_c$  dimensions).

The critical dimension itself is characterised by multiplicative logarithmic corrections, so that equations (1.2) and (1.4) become

$$c_\infty(t) \sim t^{-\alpha} |\ln t|^{\hat{\alpha}}, \quad \xi_\infty(t) \sim t^{-\nu} |\ln t|^{\hat{\nu}}, \quad (1.6)$$

and

$$\xi_L(0) \sim L(\ln L)^{\hat{\varphi}}, \quad (1.7)$$

at  $d = d_c$ , respectively. Here, we also show that the logarithmic analogue of the hyperscaling relation at the upper critical dimension is

$$\hat{\alpha} = d(\hat{\varphi} - \hat{\nu}). \quad (1.8)$$

Caution: this last scaling relation has been shown to hold *at* the upper critical dimension in a variety of models including the Ising and  $O(N)$   $\phi^4$  models, their counterparts with long-range interactions,  $m$ -component spin glasses, the percolation and Yang-Lee edge problems. It does *not* hold in *some* cases of logarithmic corrections *below* the upper critical dimension when the leading exponent  $\alpha$  vanishes. In these anomalous circumstances, an extra multiplicative logarithmic correction appears, as explained below and in reference [6, 7]. E.g., the pure Ising model in two dimensions has  $\hat{\varphi} = \hat{\nu} = 0$  but  $\hat{\alpha} = 1$ . The random-site or random-bond Ising model in  $d = 2$  has  $\hat{\varphi} = 0$ ,  $\hat{\nu} = 1/2$  but  $\hat{\alpha} = 0$ . The reason for an extra logarithm in these cases is well understood and briefly presented in section 2. Here, we are only concerned with  $d \geq d_c$ , and refer the reader to references [6, 7] for details of these anomalous cases in  $d < d_c$  dimensions.

The  $d > d_c$  hyperscaling relation (1.5) was derived in reference [4], while its logarithmic counterpart (1.8) was developed in references [6, 7]. Next, both of these derivations are summarised.

## 2. Derivation of new hyperscaling relations

We begin with more general forms for the scaling of susceptibility and correlation length in infinite volume, encompassing the leading behaviour both at and above the upper critical dimension in the thermodynamic limit, namely

$$\chi_\infty \sim t^{-\gamma} (\ln t)^{\hat{\gamma}}, \quad \xi_\infty \sim t^{-\nu} (\ln t)^{\hat{\nu}}. \quad (2.1)$$

The derivation which we are about to present involves a type of self-consistency analysis using the zeros of the partition function. The Lee-Yang zeros are those points in the complex  $h$ -plane at which the partition function  $Z_L(t, h)$  vanishes [14, 15]. Under very general conditions, Lee and Yang proved these to be located on the imaginary  $h$ -axis, although this is not a pre-requisite for what is to follow here. What is required, however, is the notion of the so-called Yang-Lee edge. In the infinite-volume limit, this is the end point of the distribution of zeros which lies closest to the real  $h$ -axis. As such, it most strongly affects critical behaviour. In line with the above ansätze, we assume that the Yang-Lee edge scales as

$$h_{\text{YL}}(t) \sim t^\Delta (\ln t)^{\hat{\Delta}}. \quad (2.2)$$

Here,  $\Delta$  is the gap exponent and  $\hat{\Delta}$  is its logarithmic counterpart. These are given through static scaling relations [6, 7]

$$\alpha = 2 + \gamma - 2\Delta, \quad \hat{\alpha} = \hat{\gamma} + 2\hat{\Delta}. \quad (2.3)$$

Our final ingredient is to promote equation (1.7) to a more general form,

$$\xi_L(0) \sim L^\vartheta (\ln L)^{\hat{\vartheta}}. \quad (2.4)$$

In each equation (2.1)–(2.4), the hatted exponents vanish above the upper critical dimension [16]. They play an important role *at*  $d_c$  itself. Circumstances in which they are non-vanishing below the upper critical dimension are not our main concern here.

We write the finite-size partition function in terms of the Lee-Yang zeros  $h_j$  as

$$Z_L(t, h) = A \prod_{j=1}^{L^d} [h - h_j(t, L)]. \quad (2.5)$$

Here, the zeros  $h_j$ , which are dependent on both  $t$  and  $L$ , are ordered such that the smaller the index  $j$ , the closer the zero is to the real  $h$ -axis. In this way,  $h_1$  is the finite-size counterpart to the Yang-Lee edge. The prefactor  $A$  plays no important role in what is to come and we henceforth drop it. The reduced free energy per unit volume is

$$f_L(t, h) = -L^{-d} \ln Z_L(t, h) \sim -L^{-d} \sum_{i=1}^{L^d} \ln [h - h_j(t, L)]. \quad (2.6)$$

Differentiating twice with respect to the field delivers the (reduced) finite-size susceptibility as

$$\chi_L(t, h) \sim \frac{\partial^2 f_L(t, h)}{\partial h^2} = \frac{1}{L^d} \sum_{j=1}^{L^d} \frac{1}{[h - h_j(t, L)]^2}. \quad (2.7)$$

From now on, we set  $h = t = 0$  and drop the corresponding arguments in  $\chi_L$  and  $h_j$ . The finite-size pseudocritical susceptibility is, therefore:

$$\chi_L \sim \frac{1}{L^d} \sum_{j=1}^{L^d} \frac{1}{h_j^2(L)}. \quad (2.8)$$

Equation (2.8) constitutes the consistency check we have been looking for; it relates the finite-size susceptibility to the  $L$ -dependency of the Lee-Yang zeros.

FSS is obtained by replacing infinite-volume correlation length in equation (2.1) by the finite-size expression (2.4),  $\xi_\infty \rightarrow \xi_L$ . This means that FSS amounts to replacing

$$t \rightarrow L^{-\frac{\alpha}{\nu}} (\ln t)^{\frac{\hat{\alpha}-\hat{\vartheta}}{\nu}}. \quad (2.9)$$

The susceptibility in equation (2.1) and the edge in equation (2.2) then take the FSS forms

$$\chi_L \sim L^{\frac{\alpha\gamma}{\nu}} (\ln L)^{\hat{\gamma}-\gamma\frac{\hat{\alpha}-\hat{\vartheta}}{\nu}}, \quad h_1(L) \sim L^{-\frac{\alpha\Delta}{\nu}} (\ln L)^{\hat{\Delta}+\Delta\frac{\hat{\alpha}-\hat{\vartheta}}{\nu}}. \quad (2.10)$$

In fact, following reference [17], one expects the higher-index zeros to scale as a function of a fraction of the total number of zeros,  $j/L^d$ . This expectation allows us to promote the second formula in equation (2.10) to

$$h_j(L) \sim \left(\frac{j}{L^d}\right)^{\frac{\hat{\Delta}}{\nu d}} \left[\ln\left(\frac{j}{L^d}\right)\right]^{\hat{\Delta} + \Delta \frac{\hat{\nu} - \hat{\phi}}{\nu}}. \quad (2.11)$$

Inserting equations (2.10) and (2.11) into the consistency expression (2.8), one finds

$$L^{\frac{\hat{\nu}\gamma}{\nu}} (\ln L)^{\hat{\gamma} - \gamma \frac{\hat{\nu} - \hat{\phi}}{\nu}} \sim L^{\frac{2\hat{\nu}\Delta}{\nu} - d} (\ln L)^{-2\hat{\Delta} \frac{\hat{\nu} - \hat{\phi}}{\nu} - 2\hat{\Delta}} \sum_{j=1}^{L^d} \left(\frac{1}{j}\right)^{2\frac{\hat{\nu}\Delta}{\nu d}}. \quad (2.12)$$

This is the equation from which we will now draw the hyperscaling relations at and above  $d_c$ .

Firstly, we assume that there are, in fact, no leading logarithmic corrections, so that  $\hat{\gamma} = 0$ ,  $\hat{\nu} = 0$ ,  $\hat{\Delta} = 0$ ,  $\hat{\phi} = 0$ . This is the circumstance above the upper critical dimension as recently confirmed by Butera and Pernici [16]. Even without these logarithmic corrections, the sum on the right hand side of equation (2.12) generates an extra logarithm if  $2q\Delta = \nu d$ . In the Ising case above  $d = d_c$ , the mean-field theory gives  $\nu = 1/2$  and  $\Delta = 3/2$ . If  $\varphi = 1$  there, one has logarithmic corrections in  $d = 6$ . This is a contradiction to the results established in reference [16]. Therefore,  $\varphi$  cannot, in fact, be 1 above  $d = d_c$ .

This is an important result. It means that the finite-size correlation length is *not* comensurate with the length above  $d = d_c$ . This is contrary to explicit statements in reference [18] and in other literature. The standard form for hyperscaling (1.3) only holds when the two length scales coincide. Above the upper critical dimension, one must account for the fact that they differ. Our main result is derived by equating the leading power-laws for  $L$  in equation (2.12). Inputting the leading static scaling relation in equation (2.3) delivers the new expression for hyperscaling in equation (1.5).

The standard leading hyperscaling relation (1.3) is valid in  $d = d_c$  dimensions so that  $\nu d_c = 2 - \alpha$ . Combining this with the new expression (1.5) leads to

$$\varphi = \frac{d}{d_c}, \quad (2.13)$$

for  $d \geq d_c$ . Since equation (1.3) is also valid in  $d < d_c$  dimensions, one gets, from equation (1.5), that  $\varphi = 1$  there. The result (2.13) for  $d \geq d_c$  agrees with an explicit analytical calculation by Brézin [19] for the large- $N$  limit of the  $N$ -vector model with periodic boundary conditions (PBC's).

However, boundary conditions do not play a role in our derivation of equation (1.5). This means, in particular, that the correlation length exceeds the actual length of the system close to  $t = 0$  for free boundary conditions (FBC's) as well as for PBC's. Again, this is contrary to many statements in the literature [18, 20–22]. In section 4, we will numerically verify that equation (2.13) indeed holds for FBC's.

The logarithmic counterpart of the hyperscaling relation comes from equating powers of logarithms in (2.12). Inserting the static relation for the correction exponents from equation (2.3), delivers equation (1.8) when  $d = d_c$ .

As stated in the Introduction, the specific heat takes on an extra logarithmic correction, beyond that coming from the hyperscaling relation (1.8), below the upper critical dimension in special circumstances. These circumstances involve the impact angle  $\phi$  at which the complex-temperature (Fisher) zeros impact onto the real axis in the thermodynamic limit. If  $\alpha = 0$ , and if this impact angle is any value other than  $\pi/4$ , an extra logarithm arises in the specific heat. This happens in  $d = 2$  dimensions for example, but not in  $d = 4$ , where  $\phi = \pi/4$  [17]. The reader is referred to reference [6, 7] for details of this anomaly.

The logarithmic critical exponent  $\hat{\phi}$  was originally introduced as  $\hat{q}$  in reference [6, 7] along with the scaling relation (1.8). That scaling relation (and its anomalous counterpart, in some  $d < d_c$  cases) was tested, and verified, against the known results in the literature there. The fact that the other exponents in equation (1.8) are universal indicates that  $\hat{\phi}$  is universal too.

Equation (1.5) is the leading-scaling counterpart of the logarithmic hyperscaling relation (1.8). Our claim in this paper is that  $\varphi$  is a new critical exponent which is universal — as are the other exponents appearing in equation (1.5). The extended hyperscaling relation (1.5) or (2.13) is, therefore, of similar

status to the other standard scaling relations. Like them, equation (1.5) holds in all dimensions, including  $d > d_c$ .

The leading part of equation (2.10) provides the leading FSS of the susceptibility and the Yang-Lee edge at the pseudocritical point  $t = 0$ . If  $\varphi$  is universal as we claim, FSS should also be universal above  $d_c$ . However, the standard belief in the literature is that above the upper critical dimension, FSS is *not* universal [18, 20–22]. In particular, until now, FSS with FBC-s has been believed to differ from FSS with periodic boundaries. We next explain how our theory differs from standard literature regarding FSS and we then go on to provide evidence which (a) identifies ours as correct and (b) explains the difference between them.

### 3. Finite-size scaling

Equation (2.10) provides the FSS of the susceptibility and the Yang-Lee edge at the pseudocritical point  $t = 0$ . In these formulae,  $\gamma$ ,  $\nu$  and  $\Delta$  assume their mean-field values 1, 1/2 and 3/2, respectively, for  $d \geq d_c$ .

Setting  $\varphi = 1$  provides the FSS at the critical dimension. For the  $O(N)$  model with  $\hat{\gamma} = (n + 2)/(n + 8)$ ,  $\hat{\nu} = (n + 2)/2(n + 8)$ ,  $\hat{\Delta} = (1 - n)/(n + 8)$  and  $\hat{\varphi} = 1/4$  [19, 23–25], one obtains

$$\chi_L(0) \sim L^2 (\ln L)^{1/2}, \quad h_1(L) \sim L^{-3} (\ln L)^{1/4}, \quad (3.1)$$

independent of  $n$ , a result already derived in reference [24] and verified numerically in the periodic Ising case in references [21, 23, 26].

Above the upper critical dimension, where there are no leading multiplicative logarithmic corrections [16], FSS for the susceptibility and edge is given by

$$\chi_L(0) \sim L^{\frac{\varphi\gamma}{\nu}}, \quad h_1(L) \sim L^{\frac{\varphi\Delta}{\nu}}, \quad (3.2)$$

respectively. If it were the case that  $\varphi = 1$ , equation (3.2) would reduce to

$$\chi_L(0) \sim L^{\frac{\gamma}{\nu}}, \quad h_1(L) \sim L^{\frac{\Delta}{\nu}}, \quad (3.3)$$

which certainly holds below  $d = d_c$  [27, 28]. Equation (3.3) also results from the Gaussian approximation. To distinguish the scaling forms (3.2) and (3.3), we refer to them as  $Q$ -FSS and Gaussian FSS, respectively [4].  $Q$ -FSS is called modified FSS in reference [29], a term earlier used in reference [30] for the adjustments to standard FSS induced by logarithmic corrections in four dimensions, following the introduction of a modified scaling variable in reference [23].  $Q$ - or modified FSS has been verified many times over for PBC's in 4,5,6,7 and 8 dimensions in references [21, 23, 24, 26, 30–37].

However, for  $\varphi$  to be a new critical exponent of similar status to the others, it must be universal. For this to be the case,  $Q$ -FSS would have to hold independent of boundary conditions. However, the standard belief for over 40 years is that systems with FBC's, in particular, have  $\chi_L \sim L^2$  or  $\varphi = 1$  [20, 22, 38, 39]. The most recent independent numerical investigation of whether Gaussian FSS or  $Q$ -FSS applies to the  $d = 5$  Ising model with FBC's, contained in reference [22], also supported the conventional belief that equation (3.3) prevails. If this were indeed the case, it would destroy the universality of  $\varphi$  and the new hyperscaling relation (1.5). In reference [4], we examined the  $d = 5$  dimensional Ising model with PBC's and FBC's. We reaffirmed that  $\varphi = d/4$  in the PBC case and we provided overwhelming evidence that  $\varphi = d/4$  in the FBC case too. Here, we focus on the latter result, since that is the one most critical for  $Q$ -theory and the most at odds with standard opinion in the literature.

### 4. Numerical evidence for universality

In this section, we provide evidence for the universality of  $\varphi$ . The evidence we present is that  $Q$ -FSS holds for FBC's, just as it does for PBC's for the Ising model in five dimensions.

As stated, it is well established that above the upper critical dimension, the Ising model on PBC lattices obeys  $Q$ -FSS at the critical point. In particular, the formula (3.2) has been verified many times for the

susceptibility at criticality [21, 31, 32, 34–37]. The corresponding expression for the Lee-Yang zeros has been verified in reference [4]. It will come as no great surprise to the reader to know that the same forms govern FSS at the pseudocritical point in the PBC case too. This has also been verified for susceptibility and the zeros in reference [4]. (In fact, Q-FSS has also been verified for the magnetization as well, both at the critical and pseudocritical points in reference [4].)

Long-standing belief is that the Q-FSS form (3.2) *does not hold* for FBC's above the upper critical dimension. Instead, standard belief is that the Gaussian form (3.3), with  $\varphi = 1$  holds there [20, 22, 38, 39]. Indeed, recent, explicit numerical support for  $\chi_L \sim L^{\varphi/\nu} = L^2$  at the critical point was given in reference [22]. We contend that this interpretation is incorrect. Our proposition is that Q-FSS applies in the FBC case above  $d = 4$ , just as it does in the PBC case. However, to observe it, one must perform FSS for FBC's at the pseudocritical point, rather than at the critical one. We will demonstrate that the infinite-volume critical point is too far away from the pseudocritical point to “feel” the finite-size scaling regime.

For the Ising model, upon which our numerical evidence is based, the partition function is


$$Z_L(t, h) = \sum_{\{s_i\}} \exp(-\beta E - hM), \quad (4.1)$$

in which  $E = -\sum_{\langle i,j \rangle} s_i s_j$ ,  $M = \sum_i s_i$ ,  $\beta = 1/kT$ ,  $h = \beta H$ ,  $H$  is the strength of an external field,  $k$  is Boltzmann's constant and  $s_i$  represents the value of an Ising spin at site  $i$  of the lattice. The summation in equation (4.1) is over spin configurations and the sum over  $\langle i, j \rangle$  is over nearest-neighbour pairs of sites on the hypercubic lattice. For a finite-size system, we measure the overall susceptibility as


$$\chi_L(t) = L^{-d} (\langle M^2 \rangle - \langle M \rangle^2). \quad (4.2)$$

For a size- $L$  hypercubic lattice, with FBC's, only  $(L - 2)^d$  sites are in the interior or bulk. Spins located on these interior sites interact with  $2d$  nearest neighbours. In this sense, they are genuinely immersed in a  $d$ -dimensional medium. The remaining  $L^d - (L - 2)^d$  spins are located on the surface of dimensionality  $d - 1$  or lower and interact directly with correspondingly fewer neighbours. These are not, therefore, fully immersed in  $d$ -dimensions. For example, with  $L = 24$  sites in each direction, the largest lattices analysed in reference [22] had only 65% of sites in the bulk. Therefore, the resulting calculations of  $\chi_L$  are not truly representative of five dimensionality. Obviously the smaller lattices are even less representative of 5D.

To get around this problem and truly probe the five-dimensionality of the FBC lattices, we decided to remove the contributions of the outer layers of sites to equation (4.2) and to other observables. We implemented this by omitting the contributions of the  $L/4$  sites near each boundary and keep only the contributions of the  $(L/2)^d$  interior sites. Each of these sites has 10 neighbours, as it should in five dimensions.



**Figure 1.** (Color online) FSS for the pseudocritical Ising susceptibility calculated on 5D lattices using all sites (upper data set, in red) and using only core sites (lower data, in blue). The upper data set appears to scale as  $\chi_L \sim L^2$ , indicative of Gaussian FSS, but this is spurious and due to the preponderance of surface sites. The lower data, which are genuinely five-dimensional, scale according to the Q-FSS form  $\chi_L \sim L^{5/2}$ , supporting the universality of  $\varphi$  (see also figure 4 (a) of reference [4]).




**Figure 2.** (Color online) FSS for the 5D Ising susceptibility at the critical point  $T_c$  from all sites (upper red data) and core sites (lower blue data). Neither Gaussian nor  $Q$ -FSS is supported (see also figure 5 (a) of reference [4]).

FSS for the susceptibility in the FBC case is represented in figure 1 where  $\chi_L$  is plotted against  $L$  on a logarithmic scale at pseudocriticality. The upper data set (red circles) corresponds to calculations of the susceptibility  $\chi_L$  in equation (4.2) using all lattice sites. The dotted line is the best fit to the form  $\chi_L \sim L^2$ . At first glance this appears to be a rather good fit, supportive of the Gaussian FSS formula (3.3) with  $\gamma/\nu = 2$  and of the traditional view of FSS for FBC lattices. However, closer inspection shows some deviation of the large- $L$  data from the dotted line. We interpret this as signaling that the apparently good fit to  $\chi_L \sim L^2$  is spurious.


The lower (blue) data set in figure 1 corresponds to calculations of  $\chi_L$  using the interior  $L/2$  lattice sites only. The dashed line is the best fit to the  $Q$ -FSS form (3.2) with  $\vartheta\gamma/\nu = 5/2$  and the line clearly describes the large- $L$  data well. This is the evidence that the Ising model defined on the five-dimensional core of the  $L^5$  lattices obeys  $Q$ -FSS (3.2) rather than Gaussian FSS (3.3). This, in turn, is the evidence for the universality of  $\vartheta$ .

In figure 2, we present FSS for the susceptibility at the critical point  $T_c$  with FBC's. Again, the upper data set represents the circumstance where all sites contribute to (4.2) and the lower data set corresponds to the usage of core sites only. Neither set of data is compatible with Gaussian or  $Q$ -FSS.

Thus  $Q$ -FSS applies at the pseudocritical point in the  $d = 5$  Ising model with FBC's, but neither  $Q$ -FSS nor Gaussian FSS applies at the critical point. To investigate why, we next examine the rounding and shifting. Both of these arise in any finite-size system because the counterpart of the divergence at  $T_c$  in infinite volume is a smoothed peak. The width of the peak is called its rounding and the location of the peak (the pseudocritical point) is shifted with respect to the critical point. If the rounding,  $\Delta T$ , is defined



**Figure 3.** (Color online) FSS of the susceptibility rounding is compatible with the  $Q$ -FSS  $\theta = \vartheta/\nu = 5/2$  rather than with the Gaussian FSS  $\theta = 1/\nu = 2$ .



**Figure 4.** (Color online) The value of the pseudocritical temperature is independent of whether it is defined from the susceptibility peak using all lattice sites or only the core (see figure 7 of reference [4]).

as the width of the susceptibility curve at half of its peak height (the “half-height width”), then one expects

$$\Delta T \sim L^{-\theta}, \quad (4.3)$$


where  $\theta$  is referred to as the rounding exponent. If  $t_L = |T_L - T_c|/T_c$  is the shift of the pseudocritical point relative to the critical one, then one has

$$t_L \sim L^{-\lambda}, \quad (4.4)$$

where  $\lambda$  is the shift exponent. For standard ( $d < d_c$ ) FSS, one normally has  $\theta = \lambda = 1/\nu$  although this is not always the case. Above the upper critical dimension this would lead to  $\theta = 2$ . For Q-FSS, one may expect that  $\theta = \lambda = \varphi/\nu = 5/2$ , but, again, this is not a requirement. Our main concern is the relative sizes of the rounding and the shifting in the FBC case. If the shifting is bigger than the rounding, then the infinite-volume critical point  $T_c$  will be too far away from the pseudocritical point to come under its effect — it will be outside the FSS regime. This would explain why FSS at the critical point is different from FSS at the pseudocritical point. In this case, FSS at  $T_c$  would certainly not be Q-FSS, but there is no reason for it to be Gaussian FSS either.


The rounding is investigated in figure 3 for the entire lattice and its core. It is obvious that the rounding is not of the standard, Gaussian type. Instead, it follows the Q-theoretic expectation, with rounding exponent  $\theta = \varphi/\nu = 5/2$ . This means that the rounding is sharper than one may naively expect, or the FSS regime is narrower than what the Gaussian theory would deliver.

In figure 4 we plot the ratio of the core susceptibility to the total susceptibility for the FBC case and for various definitions of the core. Whether the susceptibility is determined using the innermost 25%, 50% or



**Figure 5.** (Color online) The high-dimensional Ising shift exponent  $\lambda$  for FBC’s comes from the Gaussian formula  $\lambda = 1/\nu = 2$ . This means that the shifting is bigger than the rounding in the FBC case, and the critical point  $T_c$  is too far from the pseudocritical point  $T_L$  to come under its scaling effect (see also figure 6 (b) of reference [4]).





**Figure 6.** (Color online) The first two Lee-Yang zeros for Ising systems with FBC's at pseudocriticality obey Q-FSS whether the full lattice or only the core sites are used in their determination (see also figure 8 (c) of reference [4]).

75% of sites in each direction, or, indeed, whether it comprises contributions from all the sites including the boundaries, makes little difference to the location of the susceptibility peak. The shift exponent is measured in figure 5 using contributions to the susceptibility from the entire lattice. The shift exponent is clearly  $\lambda = 1/\nu = 2$ .

Thus, the shifting is indeed bigger than the rounding. Therefore, the critical point  $T_c$  is too far away from the pseudocritical point  $T_L$  to feel its effect. In other words, the finite-size susceptibility at the critical point is outside the pseudocritical FSS domain. This explains the results in figure 2 — the remoteness of  $T_c$  from  $T_L$  means that the plot is beyond the FSS regime.

To complete our investigations of equations (3.2) and (3.3) in the FBC case, the FSS for the Lee-Yang zeros is tested in figure 6 at the pseudocritical point. In fact we present the scaling of the first two zeros for FBC lattices using the contributions from all sites and from the core-lattice sites only. In each case, the zeros scale with  $L^{-q\Delta/\nu} = L^{-15/4}$  according to the Q-FSS formula on the right of equation (3.2) rather than the traditional, Gaussian formula in equation (3.3).

## 5. Dangerous irrelevant variables

The origins of the new exponent  $\varphi$  can be explained through the dangerous irrelevant mechanism in the renormalization group. Standard FSS in  $d < d_c$  may be understood by writing the finite-size free energy below the upper critical dimension as [40]

$$f_L(t, h) = b^{-d} f_{L/b}(tb^{y_t}, hb^{y_h}). \quad (5.1)$$

The correlation length is

$$\xi_L(t, h) = b \Xi_{L/b}(tb^{y_t}, hb^{y_h}), \quad (5.2)$$

which identifies  $\nu = 1/y_t$  through setting  $b = L$  and  $h = 0$ , and then taking the limit  $L \rightarrow \infty$ . At  $t = 0$ , it gives  $\xi_L \sim L$ . In the absence of the external field  $h$ , equation (5.1) can be written  $f_L(t, 0) = L^{-d} f_1[(L/\xi_\infty)^{1/\nu}, 0]$ . Thus, standard FSS is controlled by the ratio  $x = L/\xi_\infty(t)$ . Moreover, in the scaling regime where  $x \ll 1$ , the function  $f_1(x^{1/\nu}) \sim x^d$  is universal, leading to  $f_L(t) \sim t^{d\nu}$ . Differentiating this twice delivers the specific heat and standard hyperscaling (1.3).

A more complete version of equation (5.1) is [1]

$$f_L(t, h, u) = b^{-d} f_{L/b}(tb^{y_t}, hL^{y_h}, uL^{y_u}). \quad (5.3)$$

Here,  $u$  is associated with the  $\phi^4$  term in the Ginzburg-Landau-Wilson action. For  $d < d_c$ , it is a relevant scaling field, but at  $d_c$ ,  $u$  becomes marginal and above it is irrelevant. There, the Gaussian fixed point

controls the critical behaviour with  $y_t = 2$ ,  $y_h = 1 + d/2$  and  $y_u = 4 - d$  [13]. Naively differentiating equation (5.1) or (5.3) delivers functions different to those from Landau theory. This is because the limit  $u \rightarrow 0$  is singular, and should be properly accounted for. For this reason,  $u$  is termed a dangerous irrelevant variable. Its proper treatment leads to rewriting equation (5.3) as [18, 41]

$$f_L(t, h, u) = b^{-d} f_{L/b}(tb^{y_t^*}, hb^{y_h^*}) = L^{-d} f_1(tL^{y_t^*}, hL^{y_h^*}), \quad (5.4)$$

in which

$$y_t^* = y_t - \frac{y_u}{2} = \frac{d}{2}, \quad y_h^* = y_h - \frac{y_u}{4} = \frac{3d}{4}. \quad (5.5)$$

Similar considerations for the correlation length deliver

$$\xi_L(t, h, u) = L^\varphi \Xi(tL^{y_t^*}, hL^{y_h^*}). \quad (5.6)$$

In reference [18], the assumption was made that  $\varphi = 1$ . This was driven by the belief that “the correlation length  $\xi_L$  is bounded by  $L$ ” even for PBC’s. In this case, the second length scale would be needed to modify FSS [42, 43]. Introducing  $\ell_\infty(t) \sim t^{-1/y_t^*}$ , the first argument on the right-hand side of equation (5.4) or (5.6) may be written  $(\ell_\infty(t)/L)^{y_t^*}$  and  $\ell_\infty(t)$  was deemed to control FSS. It was dubbed the *thermodynamic length* by Binder [42]. Its finite-size counterpart  $\ell_L$  was called the *coherence length* in reference [29], where the so-called *characteristic length*  $\lambda_L(t)$  was also introduced as the FSS counterpart of the infinite-volume correlation length.

From our considerations, it is clear that this plethora of different lengths is unnecessary; the exponent  $\varphi$  in equation (5.6) is not bounded by 1. A direct, explicit, numerical calculation of the FSS of the correlation length for the 5D PBC model in reference [21] showed  $\xi_L \sim L^{5/4}$  there. This was verified in reference [4]. It is by now well established that the replacement of the scaling variable  $L/\xi_\infty(t)$  of standard FSS by  $L^\varphi/\xi_\infty(t)$  of  $Q$ -FSS is correct for the susceptibility, magnetization and pseudocritical point in periodic Ising models in four [21, 23], five [21, 31, 44, 45], six [32, 34, 35], seven [36] and eight [37] dimensions. The results presented here and in reference [4] support our assertion that the same holds true for FBC’s, and that  $\varphi$  is universal.

Thus, the breakdown in standard hyperscaling (1.3) above the upper critical dimension may be explained through dangerous irrelevant variables. The breakdown of FSS was less clear, however. Although the above formulation in terms of dangerous irrelevant variables does not involve explicit statements regarding boundary conditions, while it has been broadly accepted for PBC’s, Gaussian FSS was believed to hold in the FBC case. In this sense, standard FSS was not universal after all, a circumstance which was “poorly understood” [21, 46].

We have now shown that FSS for FBC’s is the same as for PBC’s at pseudocriticality, rather than at criticality and this is associated with the universality of the new exponent  $\varphi$ . However, the logarithmic counterpart to  $\varphi$  cannot be attributable to dangerous irrelevant variables, since these arise only for  $d > d_c$  and non-trivial  $\hat{\varphi}$  necessitates  $d = d_c$ . The reader is referred to reference [6, 7] for this circumstance.

## 6. Conclusions

It is well known that standard FSS is universal below the upper critical dimension  $d = d_c$  when hyperscaling holds and where the correlation length is comparable to the actual extent  $L$  of a system. Above  $d_c$ , the breakdown of standard hyperscaling is attributed to dangerous irrelevant variables in the renormalization-group approach. Although closely related to hyperscaling, FSS was so far believed to be non-universal in high dimensions, with equation (3.2) holding for FBC’s and (3.3) for PBC’s. Although this picture appeared to be supported numerically for FBC’s in references [20, 22] and for PBC’s in references [21, 23, 24, 26, 30–37], it was unexplained why the dangerous irrelevant variable mechanism should apply in one case and not in the other.

Here, we have used Lee-Yang zeros to show that the scaling mechanism is self-consistent only if the correlation length scales as a power of the length above  $d_c$ . This is the case irrespective of boundary conditions and leads to the introduction of a new scaling exponent, which we denote by  $\varphi$ . Since it is

universal,  $\varphi$  has a status similar to the critical exponents  $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\delta$ ,  $\eta$  and  $\nu$ , in notation standardised by Fisher in the 1960's. The introduction of  $\varphi$  allows one to extend the dangerous-irrelevant-variable mechanism to the correlation length through equation (5.6). FSS is then implemented by the substitution  $t \rightarrow L^{-\varphi/\nu}$ , a procedure we term “Q-FSS” to distinguish it from the standard  $t \rightarrow L^{-1/\nu}$  valid below  $d_c$ .

Here, we point out that, for the FBC lattice sizes used in reference [22], the bulk of sites are on the surface, so that the system is not genuinely five-dimensional. The resulting conclusion that  $\chi_L$  obeys Gaussian FSS is not a 5D one. For this reason, we re-examined FSS for the 5D Ising model. In order to probe the five-dimensionality of the structure, we remove contributions close to the lattice boundary. In addition to FSS at the critical temperature, we also examined pseudocriticality. Our numerical results indicate that, once the lower-dimensional effect of the peripheries is removed, the FBC lattice exhibits the *same* scaling as the PBC one at pseudocriticality, namely the one given by Q-FSS. Using the same technique at the infinite volume critical point, we find no evidence for either Gaussian FSS or for Q-FSS. Since the rounding is smaller than the shifting, we attribute this to the fact that  $T_c$  is too far from  $T_L$  to come under the effect of FSS there. This means that the conventional FSS paradigm for FBC lattices above  $d_c$  is *unsupported*, and this is particularly clear at pseudocriticality [20, 22, 38, 39]. It also offers evidence for the universality of  $\varphi$  at pseudocriticality and introduces a new, universal version of hyperscaling through equation (1.5), which is valid in all dimensions.

## Acknowledgements

We are grateful to J.-C. Walter who performed the simulations. We also thank M.E. Fisher for a number of helpful suggestions including the one to introduce the symbol  $\varphi$  for a new exponent [5]. This work is supported by the EU Programmes FP7-People-2010-IRSES (Project No. 269139) and FP7-People-2011-IRSES (Project No. 295302).

## References

1. Fisher M.E., In: Lecture Notes in Physics, **186**, Critical Phenomena, Hahne F.J.W. (Ed.), Springer, Berlin, 1983, pp. 1–139.
2. Fisher M.E., Rev. Mod. Phys., 1998, **70**, 653–681; doi:10.1103/RevModPhys.70.653.
3. Stanley H.E., Rev. Mod. Phys., 1999, **71**, S358–S366; doi:10.1103/RevModPhys.71.S358.
4. Berche B., Kenna R., Walter J.-C., Nucl. Phys. B, 2012, **865**, 115–132; doi:10.1016/j.nuclphysb.2012.07.021.
5. In reference [4], we introduced the new exponent  $q$  in analogy to  $\hat{q}$ , which was introduced in reference [6, 7] and which characterises the logarithmic-correction term there. Here we follow a suggestion made to us by M.E. Fisher who, in the 1960's standardised the notation  $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\delta$ ,  $\eta$  and  $\nu$  for the six primary critical exponents. Fisher suggested that we switch to the archaic Greek letter  $\varphi$  (“koppa”, “coppa” or “qoppa” – the source of Latin “q”) to align more closely with his, now standard, nomenclature.
6. Kenna R., Johnston D.A., Janke W., Phys. Rev. Lett., 2006, **96**, 115701; doi:10.1103/PhysRevLett.96.115701.
7. Kenna R., Johnston D.A., Janke W., Phys. Rev. Lett., 2006, **97**, 155702; doi:10.1103/PhysRevLett.97.155702.
8. Widom B., J. Chem. Phys., 1965, **43**, 3892–3897; doi:10.1063/1.1696617.
9. Widom B., J. Chem. Phys., 1965, **43**, 3898–3905; doi:10.1063/1.1696618.
10. Kadanoff L.P., Physics, 1966, **2**, 263–272.
11. Josephson B.D., Proc. Phys. Soc., 1967, **92**, 269–275; doi:10.1088/0370-1328/92/2/301.
12. Josephson B.D., Proc. Phys. Soc., 1967, **92**, 276–284; doi:10.1088/0370-1328/92/2/302.
13. Ma S.-k., Modern Theory of Critical Phenomena, Addison-Wesley, Redwood, CA, 1976.
14. Yang C.N., Lee T.D., Phys. Rev., 1952, **87**, 404–409; doi:10.1103/PhysRev.87.404.
15. Lee T.D., Yang C.N., Phys. Rev., 1952, **87**, 410–419; doi:10.1103/PhysRev.87.410.
16. Butera P., Pernici M., Phys. Rev. E, 2012, **85**, 021105; doi:10.1103/PhysRevE.85.021105.
17. Itzykson C., Pearson R.B., Zuber J.B., Nucl. Phys. B, 1983, **220**, 415–433; doi:10.1016/0550-3213(83)90499-6.
18. Binder K., Nauenberg M., Privman V., Young A.P., Phys. Rev. B, 1985, **31**, 1498–1502; doi:10.1103/PhysRevB.31.1498.
19. Brézin E., J. Phys., 1982, **43**, 15–22; doi:10.1051/jphys:0198200430101500.
20. Rudnick J., Gaspari G., Privman V., Phys. Rev. B, 1985, **32**, 7594–7596; doi:10.1103/PhysRevB.32.7594.
21. Jones J.L., Young A.P., Phys. Rev. B, 2005, **71**, 174438; doi:10.1103/PhysRevB.71.174438.

22. Lundow P.H., Markström K., Nucl. Phys. B, 2011, **845**, 120–139; doi:10.1016/j.nuclphysb.2010.12.002.
23. Kenna R., Lang C.B., Phys. Lett. B, 1991, **264**, 396–400; doi:10.1016/0370-2693(91)90367-Y.
24. Kenna R., Nucl. Phys. B., 2004, **691**, 292–304; doi:10.1016/j.nuclphysb.2004.05.012.
25. Brézin E., Le Guillou J.C., Zinn-Justin J., In: Phase Transitions and Critical Phenomena, Vol. 6, Domb C., Green M.S. (Eds.), Academic Press, New York, 1976, pp. 127-249.
26. Kenna R., Lang C.B., Nucl. Phys. B, 1993, **393**, 461–479; doi:10.1016/0550-3213(93)90068-Z.
27. Barber M.N., In: Phase Transitions and Critical Phenomena, Vol. 8, Domb C., Lebowitz J.L. (Eds.), Academic Press, New York, 1983, pp. 145-266.
28. Binder K., In: Computational Methods in Field Theory, Proceedings of the 31th Internationale Universitätswoche für Kern- und Teilchenphysik (Schladming, Austria), Gausterer H., Lang C.B. (Eds.), Springer, Berlin, 1992, pp. 59-125.
29. Brankov I.G., Danchev D.M., Tonchev N.S., Theory of Critical Phenomena in Finite-Size Systems: Scaling and Quantum Effects, World Scientific, Singapore, 2000.
30. Kenna R., Lang C.B., Nucl. Phys. B (Proc. Suppl.), 1993, **30**, 697; doi:10.1016/0920-5632(93)90305-P.
31. Aktekin N., Erkoç Ş., Kalay M., Int. J. Mod. Phys. C, 1999, **10**, 1237–1245; doi:10.1142/S0129183199001005.
32. Aktekin N., Erkoç Ş., Physica A, 2000, **284**, 206–214; doi:10.1016/S0378-4371(00)00181-3.
33. Aktekin N., J. Stat. Phys., 2001, **104**, 1397; doi:10.1023/A:1010457905088.
34. Merdan Z., Erdem R., Phys. Lett. A, 2004, **330**, 403–407; doi:10.1016/j.physleta.2004.08.030.
35. Merdan Z., Bayrih M., Appl. Math. Comput., 2005, **167**, 212–224; doi:10.1016/j.amc.2004.06.092.
36. Aktekin N., Erkoç Ş., Physica A, 2001, **290**, 123–130; doi:10.1016/S0378-4371(00)00358-7.
37. Merdan Z., Duran A., Atille D., Mulazimoglu G., Gunen A., Physica A, 2006, **366**, 265–271; doi:10.1016/j.physa.2005.10.035.
38. Watson P.G., In: Phase Transitions and Critical Phenomena, Vol. 2, Domb C., Green M.S. (Eds.), Academic, London, 1973, pp. 101–159.
39. Gunton J.D., Phys. Lett. A, 1968, **26**, 406–407; doi:10.1016/0375-9601(68)90245-4.
40. Privman V., Fisher M.E., Phys. Rev. B, 1984, **30**, 322–327; doi:10.1103/PhysRevB.30.322.
41. Brézin E., Zinn-Justin J., Nucl. Phys. B, 1985, **257**, 867–893; doi:10.1016/0550-3213(85)90379-7.
42. Binder K., Z. Phys. B, 1985, **61**, 13–23; doi:10.1007/BF01308937.
43. Binder K., Ferroelectrics, 1987, **73**, 43–67; doi:10.1080/00150198708227908.
44. Rickwardt Ch., Nielaba P., Binder K., Ann. Phys., 1994, **3**, 483–493; doi:10.1002/andp.19945060606.
45. Parisi G., Ruiz-Lorenzo J.J., Phys. Rev. B, 1996, **54**, R3698–R3702; doi:10.1103/PhysRevB.54.R3698.
46. Chen X.S., Dohm V., Phys. Rev. E, 2000, **63**, 016113; doi:10.1103/PhysRevE.63.016113.

## Новий критичний показник $\varphi$ і його логарифмічне доповнення $\hat{\varphi}$

Р. Кенна<sup>1</sup>, Б. Берш<sup>2</sup>

<sup>1</sup> Центр прикладних математичних досліджень, Університет Кавентрі, Кавентрі, Англія

<sup>2</sup> Група статистичної фізики, Інститут Жана Лямура, UMR CNRS 7198, Університет де Лоррен, 54506 Нансі, Франція

Відомо, що стандартний гіперскейлінг порушується вище верхньої критичної вимірності  $d_c$ , де критичні показники приймають класичні значення. Тут ми показуємо, що це є тому, що в стандартних формулюваннях у термодинамічній границі відстань вимірюється на масштабах кореляційної довжини. Проте, масштаб кореляційної довжини і власний масштаб довжини системи не є однакові біля чи вище вищої критичної вимірності. Вище  $d_c$  вони пов'язані алгебраїчно через новий критичний показник  $\varphi$ , тоді як біля  $d_c$  вони різняться на логарифмічні поправки, що керуються показником  $\hat{\varphi}$ . Врахування належним чином цих різних масштабів довжини дозволяє розширити гіперскейлінг до всіх вимірностей.

**Ключові слова:** гіперскейлінг, критична вимірність, кореляційна довжина, критичні індекси, співвідношення скейлінгу