

TOWARDS THE THEORY OF BRANCHING SPACES

P. I. Fomin¹ and Yu. V. Shtanov²

*Bogolyubov Institute for Theoretical Physics,
Kiev 03143, Ukraine*

We consider a spacetime of nontrivial topology formed by several pieces having common timelike boundary which plays the role of a junction between them. We establish junction conditions for fields of various spin and derive the resulting laws of wave propagation through the junction, which turn out to be quite similar for fields of all spins. As an application, we consider the case of branching four-dimensional spacetime that may arise in the context of the theory of quantum creation of a closed miniuiverse on the background of a big mother universe.

1. Introduction

In this paper, we consider the situation depicted in Fig. 1, which symbolically shows n d -dimensional Lorentzian manifolds \mathcal{M}_s , $s = 1, \dots, n$, with common $(d - 1)$ -dimensional boundary \mathcal{B} which thus plays the role of a junction between them. The boundary \mathcal{B} is assumed to be timelike, so that all the respective inner normal vector fields $n_{(s)}^a$, $s = 1, \dots, n$, at this boundary are spacelike. Our aim is to study the behavior of various fields in the space with the specified topology.




Fig. 1. Multivolume junction.

Motivation of our investigation is threefold. Firstly, it could be applied to a situation where the physical four-dimensional spacetime has a nontrivial topology that allows for branching of the type shown in Fig. 1.

¹e-mail: fomin@bitp.kiev.ua

²e-mail: shtanov@bitp.kiev.ua

Secondly, it can be applied to the study of various brane-world theories that became popular after the seminal papers [1]. In these theories, the dimensionality d of the spaces \mathcal{M}_s is usually equal to five, and the junction \mathcal{B} , which is called brane, is four-dimensional and is identified with the physical spacetime. Thus, in [2], three-brane junctions of an arbitrary number of semi-infinite four-branes were under consideration, and the whole configuration was assumed to be embedded in a six-dimensional spacetime. In this paper, however, we do not consider the space of Fig. 1 as embedded in a higher-dimensional manifold. Thirdly, in the important case $d = 2$, our investigation may be applicable to the superstring theory. Configurations of boson strings of type shown in Fig. 1 with $n = 3$ were under consideration in [3].

In brane theories, together with fields in the volume, one also considers fields whose dynamics is restricted to the brane [1]. Moreover, the action for the brane may involve the restrictions of some of the volume fields to the brane; for example, it typically involves the induced metric. However, in this paper, we restrict attention mainly to the case where the junction \mathcal{B} does not have its intrinsic dynamics and thus represents what might be called a generalization of an imaginary boundary separating two volumes in an ordinary (non-branching) space to the case where the number of volumes is greater than two.

As a concrete example of application of our theory, we consider the case of branching four-dimensional spacetime (Sec. 6.). The issue of such branching may arise in the context of the theory of quantum creation of a closed universe on the background of a big mother universe [4]. It is conceivable that the created baby universe does not become spatially separated from the mother universe, but rather remains glued with it over some common three-dimensional volume [5]. The cor-

responding situation is depicted in Fig. 2, which shows the mother universe \mathcal{M}_1 and the baby universe \mathcal{M}_2 glued over the volume \mathcal{M}_3 . All the three volume regions \mathcal{M}_1 , \mathcal{M}_2 , and \mathcal{M}_3 may evolve metrically preserving the topological configuration as shown in Fig. 2. One of the important physical questions in this situation is the issue of the behavior of various physical fields in this topology, in particular, the conditions of propagation of waves through the junction \mathcal{B} which is the common boundary of all three volume regions.

In approaching this issue, we first establish junction conditions for fields of various spins (Secs. 2., 3.) and then consider the resulting laws of wave propagation through the junction \mathcal{B} (Sec. 4.). In principle, the junction conditions at \mathcal{B} may be specified in many different ways. However, with a natural requirement that the spaces \mathcal{M}_s be treated identically, it turns out that there are precisely two versions of junction conditions for each spin. This is one of the reasons why we pay attention to spaces with the topology specified above and propose to study them in greater detail.

2. Junction conditions for the metric

We start with considering junction conditions for the metric because the actions for all other fields involve ingredients (for example, the volume element) associated with the metric. In this paper, we consider the metric g_{ab} with signature $+\dots-$. The general action for the metric can be written in the form [6]

$$S = -M^{d-2} \left(\int_{\mathcal{M}} R + 2 \sum_s \int_{\mathcal{B}} K^{(s)} \right) + \int_{\mathcal{B}} L_h, \quad (2.1)$$

where M is the Planck mass, R is the scalar curvature, and $K^{(s)}$ is the trace of the extrinsic curvature of the junction \mathcal{B} in the space \mathcal{M}_s . We impose the most natural junction conditions for the metric, namely, that the induced metric h_{ab} on \mathcal{B} is one and the same in all the spaces \mathcal{M}_s , $s = 1, \dots, n$. The Lagrangian L_h in (2.1) depends only on this induced metric.

In this paper, we use the notation and conventions of [7]. The extrinsic curvature and its trace are defined as follows:

$$K_{ab} = h^c_a \nabla_c n_b, \quad K = K_{ab} h^{ab}, \quad (2.2)$$

where $h_{ab} = g_{ab} + n_a n_b$ is the metric induced on a timelike hypersurface. The natural volume elements are implied in all the integrations over \mathcal{M} and \mathcal{B} . The cosmological-constant terms can be added to action (2.1) for each space \mathcal{M}_s , and their contribution to the resulting equations is obvious.

Variation of action (2.1) can be written in the form (see the appendix of our paper [8])

$$\delta S = - M^{d-2} \left[\int_{\mathcal{M}} G_{ab} \delta g^{ab} + \int_{\mathcal{B}} \sum_s \left(K_{ab}^{(s)} - K^{(s)} h_{ab} \right) \delta h^{ab} \right] + \int_{\mathcal{B}} \sigma_{ab} \delta h^{ab}. \quad (2.3)$$

Here, G_{ab} is the Einstein tensor, σ_{ab} is the variation of the last term in (2.1) with respect to h^{ab} , and the variation δh^{ab} is completely determined by the variation δg^{ab} of the metric in \mathcal{M} . Note that the Gibbons–Hawking boundary terms [6] in action (2.1) are required to consistently obtain the Einstein equations in the respective volume spaces without restricting variations of the metric at the junction \mathcal{B} .

Besides the metric, we may have additional fields Φ that propagate in the volume and fields φ whose dynamics is restricted to the junction. Some of the fields φ may represent restrictions of some of the volume fields Φ to the junction.

The additional set of junction conditions obtained with the account of (2.3) is then

$$\sum_s \left(K_{ab}^{(s)} - K^{(s)} h_{ab} \right) = \frac{1}{M^{d-2}} (\sigma_{ab} + \tau_{ab}), \quad (2.4)$$

where τ_{ab} is the variation of the action for the mentioned additional fields with respect to the induced metric h^{ab} .

3. Junction conditions for other fields

A complex scalar field ϕ with mass m is described by the action

$$S = \int_{\mathcal{M}} [\overline{\nabla_a \phi} \nabla^a \phi - m^2 \overline{\phi} \phi], \quad (3.1)$$

where the bar denotes complex conjugation, the integral is taken over the whole manifold $\mathcal{M} = \cup_s \mathcal{M}_s$ shown in Fig. 1, and the natural volume element is implied. The derivative $\nabla_a \phi$ may involve contribution from the gauge vector field.

In formulating the junction conditions at \mathcal{B} , we proceed from the following natural principles. Let $\phi_{(s)}$ denote the restriction of the scalar field ϕ to the space \mathcal{M}_s . We are going to relate the value of $\phi_{(s)}$ with the values of $\phi_{(r)}$, $r \neq s$ at the junction. This relation must be linear (in order to respect the superposition principle), and the spaces \mathcal{M}_s must be

regarded as physically identical. With these requirements, we arrive at the following general junction conditions at \mathcal{B} :

$$\phi_{(s)} = \alpha \sum_{r \neq s} \phi_{(r)}, \quad s = 1, \dots, n, \quad (3.2)$$

where α is some constant to be determined. Possible values of α are obtained from the additional requirement that the junction conditions (3.2) allow for non-trivial solutions at the junction. This gives only two possible values of the parameter α :

$$\alpha = \frac{1}{n-1} \quad \text{and} \quad \alpha = -1. \quad (3.3)$$

Notably, the condition $\alpha = 1/(n-1)$ simply implies the continuity of the scalar field in the space $\mathcal{M} = \cup_s \mathcal{M}_s$, i.e., the condition $\phi_{(1)} = \phi_{(2)} = \dots = \phi_{(n)}$, while the condition $\alpha = -1$ leads to the single equation $\sum_s \phi_{(s)} = 0$. To obtain other conditions at the junction, we vary the action respecting the junction conditions (3.2) and demanding that the variation be zero. General variation of action (3.1) is given by

$$\begin{aligned} \delta S = & - \int_{\mathcal{M}} [\delta \bar{\phi} (\nabla^a \nabla_a + m^2) \phi \\ & + \delta \phi (\nabla^a \nabla_a + m^2) \bar{\phi}] \\ & + \sum_s \int_{\mathcal{B}} (\delta \bar{\phi}_{(s)} \nabla_a \phi_{(s)} + \delta \phi_{(s)} \nabla_a \bar{\phi}_{(s)}) n_{(s)}^a. \end{aligned} \quad (3.4)$$

According to the value of α in (3.2), we obtain, besides the Klein–Gordon equations of motion in the volume, also the additional junction conditions.

The junction conditions for the vector field are obtained in quite a similar way. We summarize the junction conditions for the scalar and vector field in the two corresponding cases as follows:

3.1. A. Case of $\alpha = 1/(n-1)$:

$$\phi_{(p)} = \phi_{(q)}, \quad \sum_s n_{(s)}^a \nabla_a \phi_{(s)} = 0; \quad (3.5)$$

$$A_{(p)}^{\parallel a} = A_{(q)}^{\parallel a}, \quad \sum_s F_{(s)}^{ab} n_b^{(s)} = 0. \quad (3.6)$$

3.2. B. Case of $\alpha = -1$:

$$\sum_s \phi_{(s)} = 0, \quad n_{(p)}^a \nabla_a \phi_{(p)} = n_{(q)}^a \nabla_a \phi_{(q)}; \quad (3.7)$$

$$\sum_s A_{(s)}^{\parallel a} = 0, \quad F_{(p)}^{ab} n_b^{(p)} = F_{(q)}^{ab} n_b^{(q)}. \quad (3.8)$$

Here, $A^{\parallel a}$ denote the components of A^a tangent to \mathcal{B} .

Both sets of junction conditions (3.5) and (3.7) imply the sum rule for the components of the conserved current $J_a = i(\bar{\phi} \nabla_a \phi - \phi \nabla_a \bar{\phi})$ normal to the junction surface \mathcal{B} :

$$\sum_s n_{(s)}^a J_a^{(s)} = 0. \quad (3.9)$$

The junction conditions (3.5) also imply that the components of the current along the junction surface \mathcal{B} are the same in all the volume spaces.

In the case $n = 1$, i.e., where there is only one volume space with boundary, the junction conditions of type A [Eqs. (3.5), (3.6)] become the Neumann boundary conditions, while the junction conditions of type B [Eqs. (3.7), (3.8)] become the Dirichlet boundary conditions. Thus, the junction conditions obtained above can be regarded as respective generalizations of the mentioned boundary conditions to the case of $n > 1$.

Now we turn to the Dirac field. We need to relate the spinor fields $\psi_{(s)}$, $s = 1, \dots, n$, as we reach one and the same point at the junction \mathcal{B} moving in different spaces \mathcal{M}_s . The specific feature of the Dirac field is that it is referred at each point to a particular orthonormal basis (called tetrad in the case of $d = 4$). Thus, at each point $x \in \mathcal{B}$, we have to choose n orthonormal bases, one in each space \mathcal{M}_s , $s = 1, \dots, n$, to refer the corresponding values of the Dirac field to these bases. For a convenient formulation of the relations between these values, the n bases are to be chosen in a coherent way. We choose $d-1$ of the basis vectors $\{e_1^a, e_2^a, \dots, e_{d-1}^a\}$ at any point $x \in \mathcal{B}$ to be arbitrary orthonormal vectors tangent to \mathcal{B} , the same for all the spaces \mathcal{M}_s , $s = 1, \dots, n$. Then the d -th vector of the orthonormal basis in each space is determined uniquely up to sign, and we choose it to be the inner normal vector $n_{(s)}^a$, $s = 1, \dots, n$, respectively, in each of these spaces.

Next, it is clear that the spinor at one side of the junction \mathcal{B} may be related not only to the corresponding spinors themselves at the other $n-1$ sides, but also to their values in the bases reflected in the plane tangent to the junction \mathcal{B} expressed through the corresponding matrix operator of reflection $N = \gamma_{d+1} \gamma^a n_a$, where γ_{d+1} is the d -dimensional analog of the four-dimensional γ_5 matrix, which obeys the relation $N^2 = 1$.

Referring the reader to the details of the derivation presented in our paper [8], we formulate the resulting junction conditions for the Dirac field:

$$(1 \pm N) \psi_{(s)} = \frac{2}{n} \sum_r \psi_{(r)}, \quad s = 1, \dots, n. \quad (3.10)$$

In the particular case of $n = 2$, we obtain precisely the conditions which imply continuity of the Dirac field

all over \mathcal{M} . As in the previous cases, we will refer to the junction conditions (3.10) that differ in sign as to the junction conditions of type A and B, respectively, although in the case of the Dirac field, there is no qualitative difference between them.

As in the scalar case, the junction conditions (3.10) imply the sum rule (3.9) for the components of the conserved current $J^a = \bar{\psi}\gamma^a\psi$ normal to the junction.

4. Wave propagation through the junction

In this section, we apply the equations obtained to the particular interesting case of wave propagation in the space shown in Fig. 1. We shall derive the laws of wave transmission through and reflection from the junction \mathcal{B} .

First, consider the simple case of a scalar field. Let, in the region \mathcal{M}_1 , a wave that obeys the Klein-Gordon equation and propagates towards the junction \mathcal{B} be denoted by $\phi_1^{(+)}$. We denote its value at \mathcal{B} by ϕ_B and its derivative normal to the junction \mathcal{B} by $\phi'_B \equiv n_{(1)}^a \nabla_a \phi_1^{(+)}$. Then the wave which we call the reflected wave and denote by $\phi_1^{(-)}$ is constructed by imposing the same values at the junction, $\phi_1^{(-)} = \phi_B$, and by reversing the sign of the derivative normal to the junction \mathcal{B} : $n_{(1)}^a \nabla_a \phi_1^{(-)} = -\phi'_B$. For example, in the case of propagation in a flat spacetime \mathcal{M}_1 with the surface \mathcal{B} described by the equation $x_1 = 0$ in the natural spacetime coordinates $t, x \equiv (x_1, x_2, \dots, x_{d-1})$, the plane waves of this kind will be given, respectively, by $\phi_1^{(+)} = \exp(-i\omega t + ik \cdot x)$ and $\phi_1^{(-)} = \exp(-i\omega t + ik' \cdot x)$, where the wave vector k' is obtained from k by reversing its x_1 -component. The waves $\phi_s^{(-)}$, $s = 2, \dots, n$, propagating away from the junction, respectively, in the regions \mathcal{M}_s , $s = 2, \dots, n$, are constructed by imposing the boundary conditions $\phi_s^{(-)} = \phi_B$, $n_{(s)}^a \nabla_a \phi_s^{(-)} = -\phi'_B$, $s = 2, \dots, n$, at the junction \mathcal{B} . We will assume that solutions with the boundary conditions imposed exist globally in \mathcal{M}_s , $s = 1, \dots, n$, respectively.

We are looking for a solution that contains both waves falling towards \mathcal{B} and reflected from \mathcal{B} in the region \mathcal{M}_1 , but only waves propagating away from \mathcal{B} (transmitted waves) in the regions \mathcal{M}_s , $s = 2, \dots, n$. Thus, we set

$$\begin{aligned} \phi_1 &= \phi_1^{(+)} + \rho \phi_1^{(-)}, \\ \phi_s &= \tau_s \phi_s^{(-)}, \quad s = 2, \dots, n, \end{aligned} \tag{4.1}$$

where ρ is the amplitude of wave reflection and τ_s are the amplitudes of wave transmission to the spaces \mathcal{M}_s , $s = 2, \dots, n$, respectively.

To determine the amplitudes of reflection and transmission, we apply the junction conditions obtained in Sec. 3. In the case of the junction conditions (3.5), we obtain the system

$$\begin{aligned} 1 + \rho &= \tau_2 = \tau_3 = \dots = \tau_n, \\ 1 - \rho - \sum_{s=2}^n \tau_s &= 0, \end{aligned} \tag{4.2}$$

with the solution

$$\begin{aligned} \rho &= (2 - n)/n, \\ \tau_2 = \tau_3 = \dots = \tau_n &= 2/n. \end{aligned} \tag{4.3}$$

In the case of the junction conditions (3.7), we get the system

$$\begin{aligned} 1 + \rho + \sum_{s=2}^n \tau_s &= 0, \\ 1 - \rho = -\tau_2 = -\tau_3 = \dots = -\tau_n, \end{aligned} \tag{4.4}$$

and the solution that differs from (4.3) only in the sign:

$$\begin{aligned} \rho &= (n - 2)/n, \\ \tau_2 = \tau_3 = \dots = \tau_n &= -2/n. \end{aligned} \tag{4.5}$$

We see that, in both cases, the same amount of energy (the fraction $(n-2)^2/n^2$) is reflected back to the space \mathcal{M}_1 and the same equal amount of energy (the fraction $4/n^2$) is transmitted to each of the $n - 1$ spaces \mathcal{M}_s , $s = 2, \dots, n$.

The results for the case of vector fields and for weak gravitational waves are essentially the same. For the vector field, we introduce the wave $A_a^{(+)}$ propagating towards the junction \mathcal{B} in the region \mathcal{M}_1 and construct the reflected wave $A_a^{(-)}$ by keeping the component A_a^{\parallel} tangent to \mathcal{B} intact and by reversing the sign of the value of $F^{ab}n_b$ at \mathcal{B} . For a weak gravitational wave, we introduce the similar field $\delta g_{ab}^{(+)}$ and construct the corresponding reflected wave $\delta g_{ab}^{(-)}$ by keeping the perturbation of the induced metric δh_{ab} at \mathcal{B} intact and by reversing the sign of the perturbation of the extrinsic curvature δK_{ab} at \mathcal{B} . Then, proceeding in precisely the same way as we did in the scalar case, we obtain the same amplitudes of reflection and transmission. Again, the only difference between cases A and B of Sec. 3. for the vector field is in the relative phases (signs of the amplitudes) with which the waves are reflected and transmitted. The reflection and transmission amplitudes for gravitational waves will be given by (4.3).

The case of propagation of the Dirac field is also considered quite similarly to the scalar case. We denote by $\psi_1^{(+)}$ the wave that propagates towards the junction \mathcal{B} in the region \mathcal{M}_1 , and by ψ_B we denote its value at \mathcal{B} . Then the reflected wave $\psi_1^{(-)}$ is constructed by imposing the reflection boundary condition

$\psi_1^{(-)} = N\psi_B$ at \mathcal{B} and by subsequently solving the Dirac equation in \mathcal{M}_1 . Similarly, the waves that propagate away from \mathcal{B} in the spaces \mathcal{M}_s , $s = 2, \dots, n$, are constructed by imposing the conditions $\psi_s^{(-)} = N\psi_B$ at the junction \mathcal{B} and by subsequently solving the Dirac equation in \mathcal{M}_s . We assume that such solutions exist globally in \mathcal{M}_s , $s = 1, \dots, n$, as will be the case in a flat spacetime with a flat junction hypersurface considered while discussing the scalar case. With waves thus constructed, we set

$$\begin{aligned} \psi_1 &= \psi_1^{(+)} + \rho\psi_1^{(-)}, \\ \psi_s &= \tau_s\psi_s^{(-)}, \quad s = 2, \dots, n. \end{aligned} \quad (4.6)$$

Again, ρ is the coefficient of reflection, and τ_s , $s = 2, \dots, n$, are the corresponding coefficients of transmission of waves. With the upper sign in (3.10), we obtain precisely the set of equations (4.3) while, with the lower sign in (3.10), we get precisely the system of equations (4.5). Thus, we conclude that the laws of wave reflection from and transmission through the junction \mathcal{B} are similar for all the spins considered.

5. Green functions in flat branching spaces

In this section, we consider the simple case of flat spaces \mathcal{M}_s with flat common timelike boundary \mathcal{B} (see Fig. 1). We obtain the expressions for the Green functions in such a space.

In each space \mathcal{M}_s , $s = 1, \dots, n$, we can choose natural coordinates $x = \{x_1, \dots, x_d\}$ in such a way that the junction \mathcal{B} is the boundary surface $x_d = 0$ of the volume $x_d \geq 0$, and the coordinates x_1, \dots, x_{d-1} in the spaces \mathcal{M}_s are naturally identified at the junction. Let $G(x, x') \equiv D(x_1 - x'_1, \dots, x_d - x'_d)$ be any Green function (retarded, advanced, causal, etc.) for the scalar field in the Minkowski space. Then, the corresponding Green function in the space $\mathcal{M} = \cup_s \mathcal{M}_s$ is easily constructed by the method of images. We introduce the function $\tilde{G}(x, x') \equiv D(x_1 - x'_1, \dots, x_d + x'_d)$. Then the Green function $G_{\mathcal{M}}(x, x')$ in the space \mathcal{M} is given by

$$G_{\mathcal{M}}(x, x') = \begin{cases} G(x, x') \pm \frac{2-n}{n} \tilde{G}(x, x'), & x \sim x', \\ \pm \frac{2}{n} \tilde{G}(x, x'), & x \not\sim x', \end{cases} \quad (5.1)$$

where the notation $x \sim x'$ means that x and x' are in one component \mathcal{M}_s , and $x \not\sim x'$ means that x and x' are in different components \mathcal{M}_s . The upper sign in (5.1) corresponds to the junction conditions (3.5), and

the lower sign corresponds to the junction conditions (3.7).

Similar relations can be obtained for the Green functions of the vector and Dirac fields. For example, in the case of the Dirac field, the Green function $G(x, x') \equiv D(x_1 - x'_1, \dots, x_d - x'_d)$ is a matrix with spinor indices. Then we introduce the function $\tilde{G}(x, x') \equiv ND(x_1 - x'_1, \dots, x_d + x'_d)$, where N is the usual matrix of reflection acting on the index corresponding to the argument x' , and, using the junction conditions (3.10), we arrive at the same form (5.1) for the Green function.

Using expressions (5.1), one easily can obtain the renormalized vacuum stress-energy tensor. It is given by the derivatives of the Hadamard function renormalized by subtracting the Hadamard function for the Minkowski space (see [9]). We obtain

$$\langle T_{ab} \rangle_A = - \langle T_{ab} \rangle_B = \frac{2-n}{n} \langle T_{ab} \rangle_A^{(n=1)}, \quad (5.2)$$

where the labels 'A' and 'B' correspond to the junction conditions of type A and B, respectively, and the expressions for $\langle T_{ab} \rangle_{A,B}^{(n=1)}$ are standard and can be found, e.g., in [9] and references therein.

The stress-energy tensor $\langle T_{ab} \rangle_A^{(n=1)}$ typically diverges as $x_d \rightarrow 0$. For example, for a massless scalar field, we have [9]

$$\langle T_{ab} \rangle_A^{(n=1)} = \frac{1}{16\pi^2 x_d^4} g_{ab}. \quad (5.3)$$

If stress-energy tensor of such form must be added to the matter side of the Einstein equation in the volume, then its presence is inconsistent with the assumption that the spacetime is flat. This constitutes a well-known problem for curved spaces and spaces with boundaries (see [9]). However, we may simply avoid this problem in the case under consideration by requiring that exactly two copies of each field are present in the theory, one with the junction conditions A, and another with the junction conditions B. Then their contributions to the renormalized stress-energy tensor will cancel each other, as is clear from Eq. (5.2). Additionally, we must also consider quantum fluctuations of the metric that are expected to result in an effective regularization of the stress-energy tensor in the junction region. This will be the subject of the future investigations.

6. Branching universe and universe with boundary

First, let us consider a universe with spatial three-dimensional topology as shown in Fig. 2. Here, we

have three spaces \mathcal{M}_1 , \mathcal{M}_2 , and \mathcal{M}_3 with topology of a three-dimensional disk bounded by the common surface \mathcal{B} that has topology of two-sphere.




Fig. 2. Branching universe.

We assume that the topology described may arise in the context of the theory of quantum creation of a closed universe on the background of a big mother universe [4]. It is conceivable that the created baby universe does not become spatially separated from the mother universe, but rather remains glued with it over some common three-dimensional volume. Then Fig. 2 can be interpreted as showing the mother universe \mathcal{M}_1 and the baby universe \mathcal{M}_2 glued over the volume \mathcal{M}_3 . All the three volume regions \mathcal{M}_1 , \mathcal{M}_2 , and \mathcal{M}_3 that have common boundary \mathcal{B} may evolve (expand or contract) preserving the topological configuration shown. One of the important physical questions in this situation is the issue of the behavior of various physical fields, in particular, of the metric, in this topology.

Consider the case where the metrics of the pieces \mathcal{M}_1 , \mathcal{M}_2 , and \mathcal{M}_3 are the usual Friedmann–Robertson–Walker metrics given by the line element

$$ds^2 = dt^2 - a^2(t) [d\chi^2 + f^2(\chi)d\Omega_2], \quad (6.1)$$

where

$$f(\chi) = \begin{cases} \sin \chi & \text{for } \kappa = 1, \\ \chi & \text{for } \kappa = 0, \\ \sinh \chi & \text{for } \kappa = -1, \end{cases} \quad (6.2)$$

$d\Omega_2$ is the line element of the unit two-spherical geometry, and the discrete parameter κ indicates the type of the spatial geometry. The time coordinates t_s , the scale factors $a_s(t_s)$, the angles χ_s , and the functions $f_s(\chi_s)$ specified by the numbers κ_s are to be introduced for each space \mathcal{M}_s , $s = 1, 2, 3$, separately.

We consider the junction conditions (2.4) for the metric field in the absence of contribution to the right-hand side. Let the position of the junction \mathcal{B} be described by the function $\chi = \chi_*(t)$ in the metric (6.1). The components of the extrinsic curvature of the junction in the part of the space $\chi \leq \chi_*$ are given by

$$K^t_t = -\frac{d}{dt} \left[\frac{a\dot{\chi}_*}{\sqrt{1 - (a\dot{\chi}_*)^2}} \right], \quad (6.3)$$

$$K^i_j = \frac{-\delta^i_j}{\sqrt{1 - (a\dot{\chi}_*)^2}} \left[\dot{a}\dot{\chi}_* + \frac{f'(\chi_*)}{af(\chi_*)} \right], \quad (6.4)$$

where $i, j = 1, 2$ and overdot denotes the time derivative.

In general, possible motions of the junction in each of the three pieces of the volume space will be determined by the junction conditions (2.4), and this is not an easy problem even in the symmetric case that we are considering now. One special situation can be analyzed in the case where the three spaces expand in a similar way so that their Hubble parameters $H \equiv \dot{a}/a$ coincide as functions of time, which can be chosen common to all three spaces. Then solutions exist for which $\dot{\chi}_* \equiv 0$ in each of the spaces, i.e., the junction expands together with the universe. Introducing the radius $r = af(\chi_*)$ of the junction, we will have the following condition:

$$\sum_s \epsilon_s \sqrt{1 - \kappa_s (r/a_s)^2} = 0, \quad (6.5)$$

where $\epsilon_s = \text{sign } f'_s(\chi_{s*})$. Note that $\epsilon_s = 1$ for hyperbolic and flat spatial geometry ($\kappa_s = -1, 0$) while, for spherical spatial geometry ($\kappa_s = 1$), $\epsilon_s = \pm 1$. Then, with the topology shown in Fig. 2, one can conclude from Eq. (6.5) that at least two of the spaces \mathcal{M}_s must have spherical spatial geometry. Let these spaces be \mathcal{M}_1 and \mathcal{M}_2 . If, moreover, we suppose that $r \ll a_1, a_2$ and $\epsilon_1 = \epsilon_2 = -1$ (the situation actually depicted in Fig. 2) then it is necessary that the third space \mathcal{M}_3 have hyperbolic spatial geometry and $r \approx \sqrt{3}a_3$. To avoid confusion, we stress that these conditions are valid only for scaling solutions under consideration (identical Hubble parameters and $\dot{\chi}_* \equiv 0$ in each of the spaces). Also note that we have not analyzed the matter content of such universes which is necessary to produce the desired solutions.

Consider now the example of a universe whose spatial section has an internal and/or external boundary (see Fig. 3). In this case, the boundary conditions (2.4) with vanishing right-hand side imply vanishing of the extrinsic curvature of the boundary. Restricting




Fig. 3. A universe with a boundary.

analysis to the simple case of spherical boundary in a spatially flat Friedmann–Robertson–Walker spacetime, we obtain the following solution of the boundary conditions:

$$\begin{aligned} a(t) &= t_0 \left(\frac{t}{t_0} \right)^q, \\ \chi(t) &= \sqrt{q(q-1)} \left(\frac{t_0}{t} \right)^{q-1}, \end{aligned} \quad (6.6)$$

where $q > 1$. The condition $q > 1$ is a consequence of the requirement that the boundary be timelike, and it implies the power-law accelerating expansion of the universe. As it follows from (6.6), the radius of the boundary $r \equiv a\chi$ in the expanding universe increases according to the linear law $r(t) = \sqrt{q(q-1)}t$. Note that solution (6.6) can describe both a space with an outer boundary (a disk) and a space with an inner boundary (a space with a hole).

7. Discussion

Spaces with topology as that shown in Fig. 1 may naturally arise in various physical contexts, namely, in the theory of four-dimensional spacetime, in the theory of brane worlds, and in the (super)string theory. It is therefore important to study the possible junction conditions at the hypersurface \mathcal{B} and their physical consequences. In this paper, after establishing the junction conditions, we studied the issue of field propagation in spaces with the specified topology. It turns out that the laws of wave transmission through and reflection from the junction are quite similar for fields of all physical spins.

We considered the particular case of branching four-dimensional spacetime and presented a partial

solution for the metric with topology shown in Fig. 2. The aim of the subsequent investigations in this direction will be to investigate the case of branching universe in more detail and to study their physical implications. One of the ideas is to identify regions of type \mathcal{M}_3 in Fig. 2 with the observed voids (see, e.g., [10]) in the large-scale distribution of galaxies in the universe.

String configurations of type shown in Fig. 1 with $n = 3$ were studied in [3] with the natural junction conditions (3.5) for the target space coordinates on the string world sheet. It would be interesting to study such configurations in the superstring theory with the additional junction conditions (3.10) for spinor fields.

In the case of integer spin, one may wish to view the junction conditions A [with $\alpha = 1/(n-1)$] of Sec. 3. as more physical than the junction conditions B (with $\alpha = -1$) since, in the first case, the fields are continuous in the manifold \mathcal{M} while, in the second case, they are discontinuous at the surface \mathcal{B} . However, one should not discard the junction conditions of type B altogether before studying them in greater detail. This is supported by our example of flat branching spaces in Sec. 5., which shows that the presence of fields with both types of junction conditions may lead to cancellation of certain divergences in the vacuum stress-energy tensor.

Acknowledgement

The authors acknowledge useful discussion with V. P. Frolov and V. D. Gladush.

References

- [1] L. Randall and R. Sundrum, *Phys. Rev. Lett.* **83**, 3370 (1999), *E-prepr. hep-ph/9905221*; L. Randall and R. Sundrum, *Phys. Rev. Lett.* **83**, 4690 (1999), *E-prepr. hep-th/9906064*
- [2] Cs. Csáki and Yu. Shirman, *Phys. Rev. D.* **61**, 024008 (2000), *E-prepr. hep-th/9908186*
- [3] X. Artru, *Nucl. Phys. B.* **85**, 442 (1975); P. A. Collins, J. F. L. Hopkinson, and R. W. Tucker, *Nucl. Phys. B.* **100**, 157 (1975); K. Sundermeyer and A. de la Torre, *Phys. Rev. D.* **15**, 1745 (1977)
- [4] P. I. Fomin, Preprint ITP-73-137P, Kiev (1973); P. I. Fomin, *Dokl. Akad. Nauk Ukrainskoi SSR. Ser. A*, No. 9, 831 (1975)
- [5] P. I. Fomin, Yu. V. Shtanov, and O. V. Barabash, *Dopovidi Natsional'noi Akad. Nauk Ukrainy*, No. 10, 80 (2000)
- [6] G. W. Gibbons and S. W. Hawking, *Phys. Rev. D.* **15**, 2752 (1977)

- [7] R. M. Wald, General Relativity, The University of Chicago Press, Chicago (1984)
- [8] P. I. Fomin and Yu. V. Shtanov, *E-prepr. hep-th/0008183*
- [9] N. D. Birrell and P. C. W. Davies, Quantum Fields in Curved Space, Cambridge University Press, Cambridge (1982)
- [10] P. J. E. Peebles, Principles of Physical Cosmology, Princeton University Press, Princeton, New Jersey (1993)

К ТЕОРИИ ВЕТВЯЩИХСЯ ПРОСТРАНСТВ

П. И. Фомин, Ю. В. Штанов

Рассматривается пространство-время с нетривиальной топологией, состоящее из нескольких кусков, имеющих общую времениподобную границу, которая играет роль соединения между ними. Устанавливаются условия шивки для полей различных спинов и выводятся законы распространения волн через соединительную границу, которые оказываются одинаковыми для полей всех спинов. В качестве приложения рассматривается разветвленное четырехмерное пространство-время, которое может возникать в теории квантового рождения замкнутой минивселенной на фоне большой материнской вселенной.

ДО ТЕОРІЇ РОЗГАЛУЖЕНИХ ПРОСТОРІВ

П. І. Фомін, Ю. В. Штанов

Розглядається час-простір з нетривіальною топологією, який складається з кількох шматків, що мають спільну часоподібну межу, з'єднуючу їх. Знаходяться умови шивки для полів з різними спінами та закони розповсюдження хвиль крізь межу, які виявляються однаковими для усіх спінів. Як приклад застосування розглядається розгалужений чотиривимірний час-простір, який може виникати у теорії квантового народження замкнутого мінівсесвіту на фоні великого материнського Всесвіту.