

UDC 517.5

R. Y. Denis (Univ. Gorakhpur, India),
S. N. Singh, S. P. Singh (T.D.P.G. College, Jaunpur, India)

ON TRANSFORMATION FORMULAE FOR THETA HYPERGEOMETRIC FUNCTIONS

ПРО ФОРМУЛИ ПЕРЕТВОРЕННЯ ДЛЯ ГІПЕРГЕОМЕТРИЧНИХ ТЕТА-ФУНКЦІЙ

Using an identity and certain summation formulas for truncated theta hypergeometric series, we establish transformation formulas for finite bilateral theta hypergeometric series.

За допомогою однієї тотожності та формул підсумовування скорочених гіпергеометричних тета-рядів встановлено формули перетворення для скінчених двосторонніх гіпергеометричних тета-рядів.

1. Introduction, notations and definitions. Elliptic hypergeometric series and their extensions to theta hypergeometric series has become an increasingly active area of research these days. In the present paper, we have established transformation formulae for bilateral theta hypergeometric series. Special cases of the results established in this paper have also been deduced.

A modified Jacobi's theta function with argument x and nome p is defined by

$$\theta(x; p) = [x; p]_\infty [p/x; p]_\infty \equiv [x, p/x; p]_\infty. \quad (1.1)$$

Also

$$\theta(x_1, x_2, \dots, x_r; p) = \theta(x_1; p)\theta(x_2; p)\dots\theta(x_r; p)$$

and

$$[x; p]_\infty = \prod_{r=0}^{\infty} (1 - xp^r).$$

Following Gasper and Rahman [1] (Chapter 11, (11.2.5) and (11.2.53)) theta shifted factorial is defined by

$$[a; p, q]_n = \begin{cases} \theta(a; p)\theta(aq; p)\dots\theta(aq^{n-1}; p), & n > 0, \\ 1, & n = 0. \end{cases}$$

Also

$$[a; q, p]_{-n} = \frac{q^{n(n+1)/2}}{(-a)^n [q/a; q, p]_n}, \quad n \geq 1, \quad (1.2)$$

and

$$[a_1, a_2, \dots, a_r; q, p]_n = [a_1; q, p]_n [a_2; q, p]_n \dots [a_r; q, p]_n. \quad (1.3)$$

Corresponding to Spiridonov [2], theta hypergeometric series is defined by

$${}_{r+1}E_r \left[\begin{matrix} a_1, a_2, \dots, a_{r+1}; q, p; z \\ b_1, b_2, \dots, b_r \end{matrix} \right] = \sum_{n=0}^{\infty} \frac{[a_1, a_2, \dots, a_{r+1}; q, p]_n z^n}{[q, b_1, b_2, \dots, b_r; q, p]_n}, \quad (1.4)$$

where $\max\{|z|, |q|, |p|\} < 1$.

Corresponding to Spiridonov [3] a very well-poised theta hypergeometric series is defined by

$$\begin{aligned} {}_{r+1}V_r[a_1; a_6, \dots, a_{r+1}; q, p; z] &= \\ &= \sum_{n=0}^{\infty} \frac{\theta(a_1 q^{2n}; p) [a_1, a_6, \dots, a_{r+1}; q, p]_n}{\theta(a_1; p) [a_1 q/a_6, \dots, a_1 q/a_{r+1}; q, p]_n} (zq)^n = \\ &= {}_{r+1}F_r \left[\begin{matrix} a_1, q\sqrt{a_1}, -q\sqrt{a_1}, q\sqrt{a_1/p}, -q\sqrt{a_1p}, a_6, \dots, a_{r+1}; q, p; -z \\ \sqrt{a_1}, -\sqrt{a_1}, \sqrt{a_1p}, -\sqrt{a_1/p}, a_1 q/a_6, \dots, a_1 q/a_{r+1} \end{matrix} \right]. \end{aligned} \quad (1.5)$$

A truncated very well-poised theta hypergeometric series is defined by

$${}_{r+1}V_r[a_1; a_6, \dots, a_{r+1}; q, p; z]_N = \sum_{n=0}^N \frac{\theta(a_1 q^{2n}; p) [a_1, a_6, \dots, a_{r+1}; q, p]_n}{\theta(a_1; p) [a_1 q/a_6, \dots, a_1 q/a_{r+1}; q, p]_n} (zq)^n. \quad (1.6)$$

We call a series of the form

$$\sum_{k=-m}^n \frac{[a_1, a_2, \dots, a_{r+1}; q, p]_k z^k}{[q, b_1, b_2, \dots, b_r; q, p]_k}$$

a finite bilateral theta hypergeometric series.

We shall make use of the following identity:

$$\sum_{k=-m}^n \lambda_{k+m} \sum_{j=0}^{n-k} A_j = \sum_{k=-m}^n A_{k+m} \sum_{j=0}^{n-k} \lambda_j. \quad (1.7)$$

Proof of (1.7). In order to prove (1.7) let us consider the following well known identity:

$$\sum_{k=0}^n \lambda_k \sum_{j=0}^{n-k} A_j = \sum_{k=0}^n A_k \sum_{j=0}^{n-k} \lambda_j \quad (1.8)$$

(cf. Gasper, Rahman [1, p. 321], (11.6.18)). Taking $n + m$ for n and replacing k by $k + m$ in (1.8), we get (1.7) after some simplification.

Following summations are also needed in our analysis,

$${}_{10}V_9[a; b, c, d, e, q^{-n}; q, p] = \frac{[aq, aq/bc, aq/bd, aq/cd; q, p]_n}{[aq/b, aq/c, aq/d, aq/bcd; q, p]_n}, \quad (1.9)$$

where $bcdeq^{-n} = a^2q$ (cf. Gasper, Rahman [1, p. 321], (11.4.1)). Now, setting $e = aq^{n+1}$ in (1.9), we get

$${}_{8}V_7[a; b, c, a/bc; q, p]_n = \frac{[aq, aq/bc, bq, cq; q, p]_n}{[q, aq/b, aq/c, bcq; q, p]_n} \quad (1.10)$$

(cf. Gasper, Rahman [1, p. 322], (11.4.10)). Again, we have

$${}_{10}V_9[dp; a, b, dpq/c, cdpq^n/ab, q^{-n}; q, p^2] = \frac{[dpq, c/a, c/b, dpq/ab; q, p^2]_n}{[c/ab, dpq/a, c, dpq/b; q, p^2]_n} \quad (1.11)$$

(cf. Gasper, Rahman [1, p. 323], (11.4.11)). Now, taking $c = abq$ in (1.11), we get

$${}_8V_7[dp; a, b, dp/ab; q, p^2]_n = \frac{[dpq, aq, bq, dpq/ab; q, p^2]_n}{[q, abq, dpq/a, dpq/b; q, p^2]_n}. \quad (1.12)$$

We also have

$$\begin{aligned} & \sum_{k=0}^n \frac{\theta\{ad(rst/q)^k, (b/d)(r/q)^k, (c/d)(s/q)^k, (ad/bc)(t/q)^k; p\}}{\theta(ad, b/d, c/d, ad/bc; p)} \times \\ & \times \frac{[a; rst/q^2, p]_k [b; r, p]_k [c; s, p]_k [ad^2/bc; t, p]_k q^k}{[dq; q, p]_k [adst/bq; st/q, p]_k [adrt/cq; rt/q, p]_k [bcrs/dq; rs/q, p]_k} = \\ & = \frac{\theta(a, b, c, ad^2/bc; p) [arst/q^2; srt/q^2, p]_n}{d\theta(ad, b/d, c/d, ad/bc; p) [dq; q, p]_n [adst/bq; st/q, p]_n} \times \\ & \times \frac{[br; r, p]_n [cs; s, p]_n [ad^2t/bc; t, p]_n}{[adrt/cq; rt/q, p]_n [bcrs/dq; rs/q, p]_n} - \\ & - \frac{\theta(d, ad/b, ad/c, bc/d; p)}{d\theta(ad, b/d, /d, ad/bc; p)} \end{aligned} \quad (1.13)$$

(cf. Gasper, Rahman [1, p. 327], (11.6.9)). Taking $d = 1$ in the above, we get

$$\begin{aligned} & \sum_{k=0}^n \frac{\theta\{a(rst/q)^k, (b)(r/q)^k, (c)(s/q)^k, (a/bc)(t/q)^k; p\}}{\theta(a, b, c, a/bc; p)} \times \\ & \times \frac{[a; rst/q^2, p]_k [b; r, p]_k [c; s, p]_k [a/bc; t, p]_k q^k}{[q; q, p]_k [ast/bq; st/q, p]_k [art/cq; rt/q, p]_k [bcrs/q; rs/q, p]_k} = \\ & = \frac{[arst/q^2; srt/q^2, p]_n [br; r, p]_n [cs; s, p]_n [at/bc; t, p]_n}{[q; q, p]_n [ast/bq; st/q, p]_n [art/cq; rt/q, p]_n [bcrs/dq; rs/q, p]_n}. \end{aligned} \quad (1.14)$$

2. Main results. In this section we shall establish our main transformations. We start by setting

$$\lambda_k = \frac{\theta(aq^{2k}; p) [a, b, c, a/bc; q, p]_k q^k}{\theta(a; p) [q, aq/b, aq/c, bcq; q, p]_k}$$

and

$$A_k = \frac{\theta(\alpha p_1 q_1^{2k}; p_1^2) [\alpha p_1, \beta, \gamma, \alpha p_1/\beta\gamma; q_1, p_1^2]_k q_1^k}{\theta(\alpha p_1; p_1^2) [q_1, \alpha p_1 q_1/\beta, \alpha p_1 q_1/\gamma, \beta\gamma q_1; q_1, p_1^2]_k}$$

in (1.7) and using (1.10) and (1.12), we get

$$\begin{aligned} & \frac{\theta(aq^{2m}; p) [a, b, c, a/bc; q, p]_m [\alpha p_1 q_1, \alpha p_1 q_1/\beta\gamma, \beta q_1, \gamma q_1; q_1, p_1^2]_n q^m}{\theta(\alpha; p) [q, aq/b, aq/c, bcq; q, p]_m [\alpha p_1 q_1/\beta, \alpha p_1 q_1/\gamma, \beta\gamma q_1; q_1, p_1^2]_n} \times \\ & \times \sum_{k=-m}^n \frac{\theta(aq^{2m+2k}; p) [aq^m, bq^m, cq^m, aq^m/bc; q, p]_k q^k}{\theta(aq^{2m}; p) [q^{1+m}, aq^{1+m}/b, aq^{1+m}/c, bcq^{1+m}; q, p]_k} \times \end{aligned}$$

$$\begin{aligned}
& \times \frac{[q_1^{-n}, \beta q_1^{-n}/\alpha p_1, \gamma q_1^{-n}/\alpha p_1, q_1^{-n}/\beta\gamma; q_1, p_1^2]_k}{[q_1^{-n}/\alpha p_1, q_1^{-n}/\beta, q_1^{-n}/\gamma, \beta\gamma q_1^{-n}/\alpha p_1; q_1, p_1^2]_k} = \\
& = \frac{\theta(\alpha p_1 q_1^{2m}; p_1^2) [\alpha p_1, \beta, \gamma, \alpha p_1/\beta\gamma; q_1, p_1^2]_m [aq, bq, cq, aq/bc; q, p]_n q_1^m}{\theta(\alpha p_1; p_1^2) [q_1, \alpha p_1 q_1/\beta, \alpha p_1 q_1/\gamma, \beta\gamma q_1; q_1, p_1^2]_m [q, aq/b, aq/c, bcq; q, p]_n} \times \\
& \times \sum_{k=-m}^n \frac{\theta(\alpha p_1 q_1^{2m+2k}; p_1^2) [\alpha p_1 q_1^m, \beta q_1^m, \gamma q_1^m, \alpha p_1 q_1^m/\beta\gamma; q_1, p_1^2]_k q_1^k}{\theta(\alpha p_1 q_1^{2m}; p_1^2) [q_1^{1+m}, \alpha p_1 q_1^{1+m}/\beta, \alpha p_1 q_1^{1+m}/\gamma, \beta\gamma q_1^{1+m}; q_1, p_1^2]_k} \times \\
& \times \frac{[q^{-n}, bq^{-n}/a, cq^{-n}/a, q^{-n}/bc; q, p]_k}{[q^{-n}/a, q^{-n}/b, q^{-n}/c, bcq^{-n}/a; q, p]_k}. \tag{2.1}
\end{aligned}$$

Next, putting

$$\lambda_k = \frac{\theta(aq^{2k}; p) [a, b, c, a/bc; q, p]_k q^k}{\theta(a; p) [q, aq/b, aq/c, bcq; q, p]_k}$$

and

$$A_k = \frac{\theta\{\alpha(rst/q_1)^k, \beta(r/q_1)^k, \gamma(s/q_1)^k, (\alpha/\beta\gamma)(t/q_1)^k; p_1\}}{\theta(\alpha, \beta, \gamma, \alpha/\beta\gamma; p_1)} \times$$

$$\times \frac{[\alpha; rst/q_1^2, p_1]_k [\beta; r, p_1]_k [\gamma; s, p_1]_k [\alpha/\beta\gamma; t, p_1]_k q_1^k}{[q_1; q_1, p_1]_k [\alpha st/\beta q_1; st/q_1, p_1]_k [\alpha rt/\gamma q_1; rt/q_1, p_1]_k [\beta\gamma rs/q_1; rs/q_1, p_1]_k}$$

in (1.7) and using (1.10) and (1.14), we get

$$\begin{aligned}
& \frac{\theta(aq^{2m}; p) [a, b, c, a/bc; q, p]_m [\alpha rst/q_1^2; rst/q_1^2, p_1]_n [\beta r; r, p_1]_n}{\theta(a; p) [q, aq/b, aq/c, bcq; q, p]_m [q_1; q_1, p_1]_n [\alpha st/\beta q_1; st/q_1, p_1]_n} \times \\
& \times \frac{[\gamma s; s, p_1]_n [\alpha t/\beta\gamma; t, p_1]_n q^m}{[\alpha rt/\gamma q_1; rt/q_1, p_1]_n [\beta\gamma rs/q_1; rs/q_1, p_1]_n} \times \\
& \times \sum_{k=-m}^n \frac{\theta(aq^{2m+2k}; p) [aq^m, bq^m, cq^m, aq^m/bc; q, p]_k q^k}{\theta(aq^{2m}; p) [q^{1+m}, aq^{1+m}/b, aq^{1+m}/c, bcq^{1+m}; q, p]_k} \times \\
& \times \frac{[q_1^{-n}; q_1, p_1]_k [\beta(st/q_1)^{-n}/\alpha; st/q_1, p_1]_k}{[(rst/q_1^2)^{-n}/\alpha; rst/q_1^2, p_1]_k [r^{-n}/\beta; r, p_1]_k} \times \\
& \times \frac{[\gamma(rt/q_1)^{-n}/\alpha; rt/q_1, p_1]_k [(rs/q_1)^{-n}/\beta\gamma; rs/q_1, p_1]_k}{[s^{-n}/\gamma; s, p_1]_k [\beta\gamma t^{-n}/\alpha; t, p_1]_k} = \\
& = \frac{[\alpha; rst/q_1^2, p_1]_m [\beta; r, p_1]_m [\gamma; s, p_1]_m [\alpha/\beta\gamma; t, p_1]_m q_1^m}{[\alpha st/\beta q_1; st/q_1, p_1]_n [q_1; q_1, p_1]_m [\alpha rt/\gamma q_1; rt/q_1, p_1]_m [\beta\gamma rs/q_1; rs/q_1, p_1]_m} \times \\
& \times \frac{[aq, aq/bc, bq, cq; q, p]_n}{[q, aq/b, aq/c, bcq; q, p]_n} \times \\
& \times \sum_{k=-m}^n \frac{\theta\{\alpha(rst/q_1)^{k+m}, \beta(r/q_1)^{k+m}, \gamma(s/q_1)^{k+m}, \alpha(t/q_1)^{k+m}/\beta\gamma; p_1\}}{\theta(\alpha, \beta, \gamma, \alpha/\beta\gamma; p_1)} \times
\end{aligned}$$

$$\begin{aligned} & \times \frac{[\alpha(rst/q_1^2)^m; rst/q_1^2, p_1]_k [\beta r^m; r, p_1]_k [\gamma s^m; s, p_1]_k}{[q_1^{1+m}; q_1, p_1]_k [\alpha(st/q_1)^{1+m}/\beta; st/q_1, p_1]_k [\alpha(rt/q_1)^{m+1}/\gamma; rt/q_1, p_1]_k} \times \\ & \times \frac{[\alpha t^m/\beta\gamma; t, p_1]_k [q^{-n}, bq^{-n}/a, cq^{-n}/a, q^{-n}/bc; q, p]_k}{[\beta\gamma(rs/q_1)^{m+1}; rs/q_1, p_1]_k [q^{-n}/a, q^{-n}/b, q^{-n}/c, bcq^{-n}/a; q, p]_k}. \end{aligned} \quad (2.2)$$

Next, if we put

$$\lambda_k = \frac{\theta(apq^{2k}; p^2) [ap, b, c, ap/bc; q, p^2]_k q^k}{\theta(ap; p^2) [q, apq/b, apq/c, bcq; q, p^2]_k}$$

and

$$A_k = \frac{\theta\{\alpha(rst/q_1)^k, \beta(r/q_1)^k, \gamma(s/q_1)^k, \alpha(t/q_1)^k/\beta\gamma; p_1\}}{\theta(\alpha, \beta, \gamma, \alpha/\beta\gamma; p_1)} \times$$

$$\times \frac{[\alpha; rst/q_1^2, p_1]_k [\beta; r, p_1]_k [\gamma; s, p_1]_k [\alpha/\beta\gamma; t, p_1]_k q_1^k}{[q_1; q_1, p_1]_k [\alpha st/\beta q_1; st/q_1, p_1]_k [\alpha rt/\gamma q_1; rt/q_1, p_1]_k [\beta\gamma rs/q_1; rs/q_1, p_1]_k}$$

in (1.7) and using (1.12) and (1.14), we get

$$\begin{aligned} & \frac{\theta(apq^{2m}; p^2) [ap, b, c, ap/bc; q, p^2]_m q^m}{\theta(ap; p^2) [q, apq/b, apq/c, bcq; q, p^2]_m} \times \\ & \times \frac{[\alpha rst/q_1^2; rst/q_1^2, p_1]_n [\beta r; r, p_1]_n [\gamma s; s, p_1]_n [\alpha t/\beta\gamma; t, p_1]_n}{[q_1; q_1, p_1]_n [\alpha st/\beta q_1; st/q_1, p_1]_n [\alpha rt/\gamma q_1; rt/q_1, p_1]_n [\beta\gamma rs/q_1; rs/q_1, p_1]_n} \times \\ & \times \sum_{k=-m}^n \frac{\theta(apq^{2m+2k}; p^2) [apq^m, bq^m, cq^m, apq^m/bc; q, p]_k q^k}{\theta(apq^{2m}; p^2) [q^{1+m}, apq^{1+m}/b, apq^{1+m}/c, bcq^{1+m}; q, p]_k} \times \\ & \times \frac{[q_1^{-n}; q_1, p_1]_k [\beta(st/q_1)^{-n}/\alpha; st/q_1, p_1]_k [\gamma(rt/q_1)^{-n}/\alpha; rt/q_1, p_1]_k}{[(rst/q_1^2)^{-n}/\alpha; rst/q_1^2, p_1]_k [r^{-n}/\beta; r, p_1]_k [s^{-n}/\gamma; s, p_1]_k} \times \\ & \times \frac{[(rs/q_1)^{-n}/\beta\gamma; rs/q_1, p_1]_k}{[\beta\gamma t^{-n}/\alpha; t, p_1]_k} = \\ & = \frac{[\alpha; rst/q_1^2, p_1]_m [\beta; r, p_1]_m [\gamma; s, p_1]_m [\alpha/\beta\gamma; t, p_1]_m q_1^m}{[q_1; q_1, p_1]_m [\alpha st/\beta q_1; st/q_1, p_1]_m [\alpha rt/\gamma q_1; rt/q_1, p_1]_m [\beta\gamma rs/q_1; rs/q_1, p_1]_m} \times \\ & \times \frac{[aq, aq/bc, bq, cq; q, p]_n}{[q, aq/b, aq/c, bcq; q, p]_n} \times \\ & \times \sum_{k=-m}^n \frac{\theta\{\alpha(rst/q_1)^{k+m}, \beta(r/q_1)^{k+m}, \gamma(s/q_1)^{k+m}, \alpha(t/q_1)^{k+m}/\beta\gamma; p_1\}}{\theta(\alpha, \beta, \gamma, \alpha/\beta\gamma; p_1)} \times \\ & \times \frac{[\alpha(rst/q_1^2)^m; rst/q_1^2, p_1]_k [\beta r^m; r, p_1]_k [\gamma s^m; s, p_1]_k}{[q_1^{1+m}, q_1, p_1]_k [\alpha(st/q_1)^{1+m}/\beta; st/q_1, p_1]_k [\alpha(rt/q_1)^{1+m}/\gamma; rt/q_1, p_1]_k} \times \\ & \times \frac{[\alpha t^m/\beta\gamma; t, p_1]_k [q^{-n}, bq^{-n}/ap, cq^{-n}/ap, q^{-n}/bc; q, p^2]_k}{[\beta\gamma(rs/q_1)^{1+m}; rs/q_1, p_1]_k [q^{-n}/ap, q^{-n}/b, q^{-n}/c, bcq^{-n}/ap; q, p]_k}. \end{aligned} \quad (2.3)$$

3. Special cases. In this section we shall deduce certain interesting special cases of our results.

If we set $r = s = t = q_1$ in (2.2), we get after some simplification

$$\begin{aligned}
& \frac{\theta(aq^{2m}; p)[a, b, c, a/bc; q, p]_m[\alpha q_1 \alpha q_1 / \beta \gamma, \beta q_1, \gamma q_1; q_1; p_1]_n q^m}{\theta(a; p)[q, aq/b, aq/c, bcq; q, p]_m[\alpha q_1 / \beta, \alpha q_1 / \gamma, \beta \gamma q_1; q_1, p_1]_n} \times \\
& \times \sum_{k=-m}^n \frac{\theta(aq^{2m+2k}; p)[aq^m, bq^m, cq^m, aq^m/bc; q, p]_k q^k}{\theta(aq^{2m}; p)[q^{1+m}, aq^{1+m}/b, aq^{1+m}/c, bcq^{1+m}; q, p]_k} \times \\
& \times \frac{[q_1^{-n}, \beta q_1^{-n}/\alpha, \gamma q_1^{-n}/\alpha, q_1^{-n}/\beta \gamma; q_1, p_1]_k}{[q_1^{-n}/\alpha, q_1^{-n}/\beta, q_1^{-n}/\gamma, \beta \gamma q_1^{-n}/\alpha; q_1, p_1]_k} = \\
& = \frac{[\alpha, \beta, \gamma, \alpha/\beta \gamma; q_1, p_1]_m q_1^m [aq, bq, cq, aq/bc; q, p]_n}{[q_1, \alpha q_1 / \beta, \alpha q_1 / \gamma, \beta \gamma q_1; q_1, p_1]_m [q, aq/b, aq/c, bcq; q, p]_n} \times \\
& \times \sum_{k=-m}^n \frac{\theta(\alpha q_1^{2m+2k}; p_1)[\alpha q_1^m, \beta q_1^m, \gamma q_1^m, \alpha q_1^m/\beta \gamma; q_1, p_1]_k q_1^k}{\theta(\alpha; p_1)[q_1^{1+m}, \alpha q_1^{1+m}/\beta, \alpha q_1^{1+m}/\gamma, \beta \gamma q_1^{1+m}; q_1, p_1]_k} \times \\
& \times \frac{[q^{-n}, bq^{-n}/a, cq^{-n}/a, q^{-n}/bc; q, p]_k}{[q^{-n}/a, q^{-n}/b, q^{-n}/c, bcq^{-n}/a; q, p]_k}. \tag{3.1}
\end{aligned}$$

Now, setting $\alpha = \beta \gamma$ in (3.1), we get

$$\sum_{k=0}^n \frac{\theta(aq^{2k}; p)[a, b, c, a/bc; q, p]_k}{\theta(a; p)[q, aq/b, aq/c, bcq; q, p]_k} = \frac{[aq, aq/bc, bq, cq; q, p]_n}{[q, aq/b, aq/c, bcq; q, p]_n}, \tag{3.2}$$

which is (1.9).

Again, if we take $\alpha = \beta \gamma q_1$ in (3.1), we get

$$\begin{aligned}
& \frac{\theta(aq^{2m}; p)\theta(\beta q_1, \gamma q_1, \beta \gamma q_1^{1+n}, q_1^{1+n}; p_1)[a, b, c, a/bc; q, p]_m q^m}{\theta(a; p)\theta(q_1, \beta \gamma q_1, \beta q_1^{1+n}, \gamma q_1^{1+n}; p_1)[q, aq/b, aq/c, bcq; q, p]_m} \times \\
& \times \sum_{k=-m}^n \frac{\theta(aq^{2m+2k}; p)[aq^m, bq^m, cq^m, aq^m/bc; q, p]_k q^k}{\theta(aq^{2m}; p)[q^{1+m}, aq^{1+m}/b, aq^{1+m}/c, bcq^{1+m}; q, p]_k} \times \\
& \times \frac{[q_1^{-n}, q_1^{-n-1}/\beta, q_1^{-n-1}/\gamma, q_1^{-n}/\beta \gamma; q_1, p_1]_k}{[q_1^{-n-1}/\beta \gamma, q_1^{-n}/\beta, q_1^{-n}/\gamma, q^{-n-1}; q_1, p_1]_k} = \\
& = \frac{[\beta, \gamma; q_1, p_1]_m q_1^m [aq, bq, cq, aq/bc; q, p]_n}{[\gamma q_1^2, \beta q_1^2; q_1, p_1]_m [q, aq/b, aq/c, bcq; q, p]_n} \times \\
& \times \sum_{k=-m}^n \frac{\theta(\beta \gamma q_1^{2m+2k+1}; p_1)[\beta q_1^m, \gamma q_1^m; q_1, p_1]_k q_1^k}{\theta(\beta \gamma; p_1)[\beta q_1^{2+m}, \gamma q_1^{2+m}; q_1, p_1]_k} \times \\
& \times \frac{[q^{-n}, bq^{-n}/a, cq^{-n}/a, q^{-n}/bc; q, p]_k}{[q^{-n}/a, q^{-n}/b, q^{-n}/c, bcq^{-n}/a; q, p]_k}. \tag{3.3}
\end{aligned}$$

Next, if we put $m = 0$ in (3.1), we get

$$\begin{aligned} & \frac{[\alpha q_1, \beta q_1, \gamma q_1, \alpha q_1/\beta\gamma; q_1, p_1]_n}{[q_1, \alpha q_1/\beta, \alpha q_1/\gamma, \beta\gamma q_1; q_1, p_1]_n} \times \\ & \times \sum_{k=0}^n \frac{\theta(aq^{2k}; p)[a, b, c, a/bc; q, p]_k [q_1^{-n}, \beta q_1^{-n}/\alpha, \gamma q_1^{-n}/\alpha, q_1^{-n}/\beta\gamma; q_1, p_1]_k q^k}{\theta(a; p)[q, aq/b, aq/c, bcq; q, p]_k [q_1^{-n}/\alpha, q_1^{-n}/\beta, q_1^{-n}/\gamma, \beta\gamma q_1^{-n}/\alpha; q_1, p_1]_k} = \\ & = \frac{[aq, bq, cq, aq/bc; q, p]_n}{[q, aq/b, aq/c, bcq; q, p]_n} \times \\ & \times \sum_{k=0}^n \frac{\theta(aq_1^{2k}; p_1)[\alpha, \beta, \gamma, \alpha/\beta\gamma; q_1, p_1]_k [q^{-n}, bq^{-n}/a, cq^{-n}/a, q^{-n}/bc; q, p]_k q_1^k}{\theta(\alpha; p_1)[q_1, \alpha q_1/\beta, \alpha q_1/\gamma, \beta\gamma q_1; q_1, p_1]_k [q^{-n}/a, q^{-n}/b, q^{-n}/c, bcq^{-n}/a; q, p]_k}. \end{aligned} \quad (3.4)$$

If we set $p = p_1 = 0$ and $q_1 = q$ in (3.4), we get the following interesting transformation:

$$\begin{aligned} {}_{10}\phi_9 \left[\begin{matrix} a, q\sqrt{a} - q\sqrt{a}, b, c, a/bc, \beta q^{-n}/\alpha, \gamma q^{-n}/\alpha, q^{-n}/\beta\gamma, q^{-n}; q; q \\ \sqrt{a}, -\sqrt{a}, aq/b, aq/c, bcq, q^{-n}/\alpha, q^{-n}/\beta, q^{-n}/\gamma, \beta\gamma q^{-n}/\alpha \end{matrix} \right] = \\ = \frac{[aq, aq/bc, bq, cq, \alpha q/\beta, \alpha q/\gamma, \beta\gamma q; q]_n}{[aq/b, aq/c, bcq, \alpha q, \beta q, \gamma q, \alpha q/\beta\gamma; q]_n} \times \\ \times {}_{10}\phi_9 \left[\begin{matrix} \alpha, q\sqrt{a}, -q\sqrt{a}, \beta, \gamma, \alpha/\beta\gamma, bq^{-n}/a, cq^{-n}/a, q^{-n}/bc, q^{-n}; q; q \\ \sqrt{a}, -\sqrt{a}, \alpha q/\beta, \alpha q/\gamma, \beta\gamma q, q^{-n}/a, q^{-n}/b, q^{-n}/c, bcq^{-n}/a \end{matrix} \right]. \end{aligned} \quad (3.5)$$

Now, letting $\beta \rightarrow 1$ in (3.5) we get the following summation of a truncated very well poised ${}_6\phi_5$

$${}_6\phi_5 \left[\begin{matrix} a, q\sqrt{a}, -q\sqrt{a}, b, c, a/bc; q; q \\ \sqrt{a}, -\sqrt{a}, aq/b, aq/c, bcq \end{matrix} \right]_n = \frac{[aq/aq/bc, bq, cq; q]_n}{[q, aq/b, aq/c, bcq; q]_n}.$$

It is evident that several other interesting results involving theta hypergeometric functions can be established.

Acknowledgement. The authors express deep appreciation to the referee for his valuable suggestions. This has certainly improved the quality of the paper. They are thankful to the Department of Science and Technology, Govt. of India, New Delhi, for support under major research projects No. SR/S4/MS-461/07 dtd.13.2.2008 entitled “A study of basic hypergeometric functions with special reference to Ramanujan mathematics”; No. SR/S4/MS:524 dtd.10.2.2008 entitled “Glimpses of Ramanujan’s mathematics in the field of q -series” and No. F.6-2(23)/2008(MRC/NRCB) dtd.5.6.2009, entitled “Investigations of Ramanujan’s work in the field of basic hypergeometric series” sanctioned to them, respectively. The first author is also thankful to the Indian Society of Mathematics and Mathematical Sciences (ISMAMS) for sponsoring his project.

1. Gasper G., Rahman M. Basic hypergeometric series. – Second ed. – Cambridge Univ. Press, 2004.
2. Spiridonov V. P. Theta hypergeometric series // Proc. NATO ASI Asympt. Combin. Appl. Math. Phys. (St.Petersburg, July 9–22, 2001). – Dordrecht: Kluwer Acad. Publ., 2002. – P. 307–327.
3. Spiridonov V. P. An elliptic incarnation of the Bailey chain // In. Math. Res. Not. – 2002. – 37. – P. 1945–1977.

Received 14.08.11,
after revision – 15.06.12