

THE LEVEL OF STEROID HORMONES IN THE BLOOD SERUM OF WOMEN WITH ENDOCRINE FORMS OF STERILITY AND NARROW PELVIS

O.V. Tyan

SUMMARY

In our research , in order to study the content of steroid hormones in the blood serum of women with various forms of endocrine infertility. Hormones assessed by the level of hormones in the blood, the size of the pelvis were determined using measurement of pelvic. The positive correlation between the concentration of steroid hormones and the degree of narrowing of the pelvis in women with infertility.

2012, 15, 2, 3 (58)

3 , - . / 2,5 -
 , 58% (29) , / 2
 20% , 22%
 29,87% / . 1

(±m)

	, /	, /	, /	, /
	0,49±0,02	12,56±0,15	42,35±0,93	682,24±39,65
	1,54±0,03*	6,55±0,71*	36,31±3,39	478,63±36,74*
	3,67±0,08*	20,7±1,99*	50,69±4,48	442,35±41,12*

:* - <0,05

18,1 (/), . . -
 : 7 , - () 15,2%.
 , 2 (1). - 1.
 50,69±4,48 / , 19,7% 2.
 , -
 35% - 3.
 , -
 (r_{xy} 0,40), 4.
 .. -
 .. (r_{xy} 0,43), . . -
 , .
 23,6. 30 () - 5.
 -
 (0,12-0,25 /), . . - (r_{xy} -0,26 r_{xy} -0,18,) -
 , , , -
 (r_{xy} 0,43), . . -
 . 1. . . -
 (r_{xy} 0,31 -), . . - /
 , - , . . // -
 , -2010.- 7.- .95-100. -
 2. . . // -
 9,47 12,25 /). -2009.- 3 (39).- .206. -
 (19,07) 3. . . -
 (r_{xy} 0,25). / . . // -
 20 2005.- .11, 4.- .35-40. -
 4. . . -
 (13,1, 15,3, 17,2 « » / . . -

, . . . , . . . // - 4.- .16-19.
 .-2000.- 1.- .43-44. 6. . . -
 5. - . . . / . . / . . , . . // -
 , . . , . . // - .-2006.- 2 (26).
 - .-2005.- .5, - .42-45.