

LASER COOLING IN AN ELECTRON STORAGE RING AND ITS LIMITS

A.N. Lebedev

P.N. Lebedev Physical Institute

119924 Leninsky Prosp. 53, Moscow, Russia

E-mail: lebedev@sci.lebedev.ru; tel. (095) 132-5582

The evolution of synchrotron and betatron emittances of an electron beam under action of laser irradiation and consequent emission of hard quanta is analyzed. Dependencies of the cooling rate on structure functions at the irradiation point as well as on the parameters of the electron and laser beams are calculated and optimized. The invariance of the sum of the decrements is proved.

The work is fulfilled within the frames of NATO grant Sfp-977982.

PACS: 41.60.Cr

1. INTRODUCTION

The possibility of generation of monochromatic X-ray radiation by backward scattering of laser light on a relativistic electron beam attracts now special attention [1,2]. The frequency transformation easily follows from simple kinematic relations and has an order of magnitude of $4\gamma^2$ where γ is Lorentz factor. Besides, the scattered X-rays are well directed (angle $\approx 1/\gamma$) what is typical for radiation from high energy electrons. A weak point of the method is a rather small cross-section which imposes serious requirements upon the laser power and the beam density.

It is rather obvious that electron beams circulating in a storage ring are preferable from this point of view if their life-time is large enough. The latter depends on many factors including transverse spreading of the beam inherent in the method and caused by the recoil of emitted hard quanta. This effect is well known for cyclic accelerators in connection with quantum fluctuations of synchrotron radiation. However, in our case a quantum is essentially harder and requirements to the beam transverse size are more stressed.

Similarly, one could count on radiation cooling also inherent in the scattering process. Really, a radiation quantum should be re-emitted practically along the instantaneous velocity of a relativistic electron which gets both longitudinal and transverse recoil momentum. The first is restored by the RF compensating system while the second produces a radiation "friction" exactly in the same way as in synchrotrons. Although one can not expect really strong damping for existing parameters the effect has to be considered because the spectral and angular distribution of the scattered light differs from that in synchrotrons and depends on parameters of the laser beam.

2. LASER COOLING

We neglect below intrinsic damping due to synchrotron radiation and consider electrons performing independent synchrotron and betatron oscillations:

$$y = x + R\psi u; \quad x = \sqrt{\varepsilon/\pi} \beta^{1/2} \cos \int \frac{ds}{\beta}; \quad (1)$$

$$x' = \sqrt{\varepsilon/\pi} \beta^{-1/2} \left(\frac{\beta'}{2} \cos \int \frac{ds}{\beta} - \sin \int \frac{ds}{\beta} \right). \quad (2)$$

Here u is a relative energy deviation from the equilibrium value, β and ψ are periodic structure functions of the magnetic system, R is the mean radius of the equilibrium orbit, primes denote derivatives with respect to the orbit arc s . The value ε has the meaning of the area enclosed by a phase trajectory ellipsis in the phase plane (x, x') being an integral of motion. For bounding phase trajectories it is identified as a transverse emittance. Being expressed via phase plane coordinates it is equal to

$$\frac{\varepsilon}{\pi} = \frac{x^2}{\beta} + \frac{1}{\beta} \left(\frac{\beta'}{2} x - \beta x' \right)^2. \quad (3)$$

Meeting a laser photon at the light and electron beams crossing point the electron energy is instantaneously changed by the value Δu keeping the coordinate y and the instantaneous velocity y' constant. The latter means that the scattered photon is emitted perfectly along the electron trajectory. With the same precision one can neglect the energy change when a relatively soft laser photon is absorbed. As a result, the integral ε experiences the instantaneous change:

$$\frac{\Delta \varepsilon}{2\pi R \Delta u} = x \frac{\psi'}{\beta} + \beta \frac{\psi}{\beta} \frac{1}{2} \frac{\psi'}{\beta} \frac{1}{2} \frac{\psi}{\beta} - \beta^{3/2} x \frac{\psi'}{\beta} \frac{1}{2} \frac{\psi}{\beta} + \dots \quad (4)$$

Here and below the structure functions and their derivatives are taken at the crossing point. A destination of the second order changes will be considered later.

To find the average variation rate $d\varepsilon/ds$ the expression above is to be multiplied by the scattering probability $P(\Delta u, y, y')$ and averaged over all betatron phases. To do this we present the probability as an expansion

$$P = P_0 + \frac{\partial P}{\partial y} y + \frac{\partial P}{\partial y'} y' + \frac{\partial P}{\partial u} u + \frac{\partial P}{\partial u'} u'. \quad (5)$$

The first term here is the probability of an "ideal" collision with the equilibrium electron, the second and the third ones describe coordinate and angular discrepancies between the electron and light beam axis. The last term corresponds to a possible influence of lack of synchronization between electron and light pulses.

Note that after averaging over phases denoted below by angular brackets all terms of the first order with re-

spect to y, y', u, u' vanish. Besides, it follows from Eqs. (1)-(2) that

$$\langle xy \rangle = \frac{\varepsilon}{2\pi} \beta; \quad \langle xy' \rangle = \frac{\varepsilon}{2\pi} \beta'; \quad (6)$$

$$\langle x'y \rangle = \frac{\varepsilon}{4\pi} \beta'; \quad \langle x'y' \rangle = \frac{\varepsilon}{2\pi\beta} (\beta'^2 / 4 + 1). \quad (7)$$

So we get

$$\left\langle \frac{\Delta \varepsilon}{\varepsilon} \right\rangle = \Delta u R \left(-\psi \frac{\partial P}{\partial y} - \psi' \frac{\partial P}{\partial y'} \right). \quad (8)$$

The same arguments give for the synchrotron emittance

$$\left\langle \frac{\Delta \varepsilon_s}{\varepsilon_s} \right\rangle = \Delta u \left(R\psi \frac{\partial P}{\partial y} + R\psi' \frac{\partial P}{\partial y'} + \frac{\partial P}{\partial u} \right). \quad (9)$$

Besides the re-emission process, the transverse emittance is influenced by the RF field necessary for radiation losses compensation. We shall suppose it being concentrated within a narrow accelerating gap normal to the equilibrium orbit. A particle gets there an instantaneous increase in energy, keeping y and u' constant. There is a simultaneous change of y' because the accelerating field does not change the transverse momentum p_y .

For this reason the change of the emittance has to be calculated under condition

$$\Delta y' = -\Delta \frac{p_y}{p} u = -y' u \quad (10)$$

as far as in the ultrarelativistic case the relative changes of energy and of total momentum P are equal to each other. Moreover, the probability of the gain is now identically equal to unity because the energy income does not depend on the possible scattering at previous turns.

Noting that $-P\Delta u$ is equal to the relative energy W emitted per one turn we get the increments of betatron and synchrotron oscillations damping:

$$\Gamma_b = -R\psi \frac{\partial W}{\partial y} - R\psi' \frac{\partial W}{\partial y'} + W; \quad (11)$$

$$\Gamma_s = R\psi \frac{\partial W}{\partial y} + R\psi' \frac{\partial W}{\partial y'} + \frac{\partial W}{\partial u}.$$

They have much common with the usual radiation damping decrements but contain the local values of the structure functions. In particular the theorem about the decrements sum [3] looks as

$$\Gamma_b + \Gamma_s = W + \frac{\partial W}{\partial u}. \quad (12)$$

It says that the coordinate and angular discrepancies of laser and electron beams do not influence the total phase volume and yield a decrement re-distribution only. By the way, the term proportional to the coordinate shift between the beams vanishes unless the beams have a zero crossing angle.

Bearing in mind that the intensity of radiation of a relativistic particle is proportional to the square of its energy the relation (12) can be rewritten as

$$\Gamma_b + \Gamma_s = 3W. \quad (13)$$

The laser cooling as opposed to synchrotron radiation one depends on the laser power and thus on the final output of hard quanta. In certain ambitious projects the damping time can be less than a millisecond. This might provide a serious limitation of the emittance growth due to quantum fluctuations discussed below.

3. EXITATION BY RECOIL MOMENTUM FLUCTUATIONS

A quantum nature of radiation is described by the next terms of expansion of ε_b and ε_s over powers of Δu . Bearing in mind that in the limit of $\hbar \rightarrow 0$

$$\langle (\Delta u)^2 P_0 \rangle = E_q W$$

where

$$E_q = \langle (\Delta u)^2 P_0 \rangle / \langle \Delta u P_0 \rangle,$$

the scattering process results in the emittances increase rate:

$$\frac{d\varepsilon_b}{ds} = \pi R^2 E_q W \left[\frac{\psi^2}{\beta} + \beta^2 \left(\frac{\psi}{\beta^{1/2}} \right)^2 \right]; \quad (14)$$

$$\frac{d\varepsilon_s}{ds} = \pi R \frac{q\alpha}{\Omega} E_q W, \quad (15)$$

where Ω is a synchrotron frequency, q is a harmonic number, and $\alpha = \bar{v}$ is a momentum compaction factor.

This rate has to be compared with damping due to laser cooling. Note that both are proportional to the laser power so that the final steady-state emittance has a universal value of order of

$$\varepsilon_b \approx \frac{1}{\Gamma_b} \frac{d\varepsilon_b}{ds} \approx \pi R^2 E_q \left[\frac{\psi^2}{\beta} + \beta^2 \left(\frac{\psi}{\beta^{1/2}} \right)^2 \right]. \quad (16)$$

This value determines, of course, whether the stored electrons can be exploited for a long time or they would be burning down and require continual reinforcement.

4. ON OPTIMIZATION OF THE STRUCTURE FUNCTIONS

It is easy to see that to make the emittance growth smaller the value of

$$U = \frac{\psi^2}{\beta} + \beta^2 \left(\frac{\psi}{\beta^{1/2}} \right)^2 \quad (17)$$

should be as small as possible at the crossing point while the mutual geometry of the electron and light beams can influence the decrements redistribution only (here and below β, ψ are again the structure functions of s). Note that they are not independent as far as $\psi(s)$ is a periodic solution of

$$\psi'' + g(s)\psi = K(s)/R \quad (18)$$

while the amplitude function $\beta(s)$ satisfies the nonlinear equation

$$\left(\beta^{1/2} \right)'' + g(s)\beta^{1/2} = \beta^{-3/2} \quad (19)$$

with the same focussing function $g(s)$. Here $K(s)$ is the equilibrium orbit curvature.

Multiplying Eq.(18) by $\beta^{1/2}$ and Eq.(19) by Ψ gives the general equation relating the structure functions

$$\beta \frac{d}{ds} \beta \frac{d}{ds} \frac{\Psi}{\beta^{1/2}} + \frac{\Psi}{\beta^{1/2}} = \frac{K}{R} \beta^{3/2}. \quad (20)$$

Using this relation and differentiating U with respect to S to find an extremum we have:

$$U' = 2 \frac{K}{R} \beta^{3/2} \left(\frac{\Psi}{\beta^{1/2}} \right)'. \quad (21)$$

Thus, U reaches its extremal values at the same points where $\Psi / \beta^{1/2}$ does while $U_{extr} = \left(\Psi / \beta^{1/2} \right)_{extr}$.

Within a straight section U is a non-zero constant which can be expressed in terms of the positive β function. Really, considering $\varphi = \int ds / \beta$ as an independent variable in Eq. (20) we have

$$\left[\frac{d^2}{d\varphi^2} + 1 \right] \left(\frac{\Psi}{\beta^{1/2}} \right) = \frac{K}{R} \beta^{3/2} \quad (22)$$

with the periodicity conditions on the interval $0 \leq \varphi \leq 2\pi\nu$ where $\nu = (2\pi)^{-1} \int_0^{2\pi R} ds / \beta$ is the betatron oscillation frequency. The solution is straightforward:

$$\frac{\Psi}{\beta^{1/2}} = \int_0^\varphi \frac{K}{R} \beta^{3/2} \sin(\varphi - \varphi') d\varphi' + \frac{1}{2 \sin \pi \nu} \int_0^{2\pi \nu} \frac{K}{R} \beta^{3/2} \sin(\pi \nu + \varphi - \varphi') d\varphi'. \quad (23)$$

At a point of an extremum where $\left(\Psi / \beta^{1/2} \right)' = 0$ the function U reaches the value

$$U_{extr} = \left| \int_0^{2\pi \nu} \frac{K}{2\pi R \sin \pi \nu} \beta^{3/2} e^{i\varphi'} d\varphi' \right|^2 \quad (24)$$

So, one can say that the excitation of oscillations always takes place. To minimize it, a defocusing lens could be helpful at the crossing point as well as negative curvature portions of the equilibrium orbit with large values of β function.

REFERENCES

1. F. Arutyunian, V. Tumanian // *Phys. Lett.* 1963, p.176; R. Milburn // *Phys. Rev. Lett.* 1963, v.10, p.75.
2. A. Agafonov et al. // *Problems of Atomic Science and Technology. Series: Nuclear Physics Investigations.* 2001, №2(37), p.126.
3. A. Kolomensky and A. Lebedev. *The Theory of Cyclic Accelerators.* N. Hol. Publ. Co. 1963.

ЛАЗЕРНОЕ ОХЛАЖДЕНИЕ В ЭЛЕКТРОННОМ НАКОПИТЕЛЬНОМ КОЛЬЦЕ И ЕГО ПРЕДЕЛЫ

А.Н. Лебедев

Исследована эволюция поперечного и продольного эмиттансов электронного пучка при взаимодействии его с лазерным излучением и последующей эмиссии жестких квантов. Рассчитаны и оптимизированы зависимости скорости охлаждения от структурных функций в точке взаимодействия, а также параметры электронного и лазерного пучков. Доказана инвариантность суммы декрементов. Найден минимальный фазовый объем, определяемый эффектом отдачи жестких квантов. Работа выполнена при поддержке гранта НАТО SfP-977982.

ЛАЗЕРНЕ ОХОЛОДЖЕННЯ В ЕЛЕКТРОННОМУ НАКОПИЧУВАЛЬНОМУ КІЛЬЦІ І ЙОГО МЕЖІ

А.М. Лебедев

Розглянуто еволюцію синхротронного і бетатронного еміттанса електронного пучка під дією лазерного опромінення і наступного випромінювання твердих квантів. Знайдені і оптимізовані залежності швидкості охолодження пучка від структурних функцій у точці зустрічі електронного і фотонного пучків і від параметрів останніх. Доведено інваріантність суми декрементів. Знайдено мінімальний фазовий об'єм, обумовлений ефектом віддачі твердих квантів. Робота виконана за підтримкою гранту НАТО SfP-977982.