

Reentrant transitions of a mixed-spin Ising model on the diced lattice*

M.Jaščur, J.Strečka

Department of Theoretical Physics and Astrophysics,
Faculty of Science, P. J. Šafárik University,
Park Angelinum 9, 040 01 Košice, Slovak Republic

Received July 18, 2005

Magnetic behaviour of a mixed spin-1/2 and spin-1 Ising model on the diced lattice is studied using an exact star-triangle mapping transformation. It is found that the uniaxial as well as biaxial single-ion anisotropy acting on the spin-1 sites may potentially cause a reentrant transition with two consecutive critical points. Contrary to this, the effect of next-nearest-neighbour interaction between the spin-1/2 sites possibly leads to a reentrant transition with three critical temperatures in addition to the one with two critical points only. The shape of the total magnetization versus temperature dependence is particularly investigated for the case of ferrimagnetically ordered system.

Key words: *reentrant transition, Ising model, exact solution, star-triangle transformation, uniaxial and biaxial single-ion anisotropy*

PACS: *75.10.Hk, 05.50.+q, 75.40*

1. Introduction

Over the last six decades, much effort has been devoted to determining the criticality and other statistical properties of various lattice-statistical models, which would enable a deeper understanding of order-disorder phenomena in solids. The planar Ising model holds a prominent place in the equilibrium statistical physics from this point of view as it represents a rare example of exactly solvable lattice-statistical model since Onsager's pioneering work [1]. It is noteworthy that an exact tractability of two-dimensional Ising systems [2] is of immense importance due to the convincing evidence provided for many controversial results predicted in the phase transition theory. Moreover, the exactly soluble models provide a good testing ground for approximative theories. It should be emphasized, however, that a precise treatment of planar Ising models is usually connected with the usage of sophisticated mathematical methods, which consequently lead to considerable difficulties when applying them to more complicated but more realistic systems.

*This work was financially supported under the grants VEGA 1/2009/05 and APVT 20-005204.

The Ising systems consisting of mixed spins of different magnitudes, the so-called *mixed-spin Ising models*, are among the most interesting extensions of the standard spin-1/2 Ising model which are currently of great research interest. These models enjoy a considerable attention mainly due to much richer critical behaviour they display compared with their single-spin counterparts. Magnetic properties of the mixed-spin Ising models can be essentially modified, for instance, by the presence of single-ion anisotropy acting on the $S \geq 1$ spins. Indeed, this effect induced by the crystal field of ligands may potentially cause a tricritical phenomenon, a change of the character of phase transition from a second-order to a first-order one. Another aspect which started to attract an appreciable interest towards the mixed-spin Ising models can be related to a theoretical modelling of magnetic structures suitable for describing a ferrimagnetism of certain class of insulating materials. Accordingly, the ferrimagnetic mixed-spin Ising models are also very interesting from the experimental viewpoint, first of all in connection with many possible technological applications of ferrimagnets (e.g. thermomagnetic recording).

Despite the intensive studies, there are only a few examples of exactly solvable mixed-spin models so far. Using the generalized form of decoration-iteration and star-triangle transformations, the mixed spin-1/2 and spin- S ($S \geq 1$) Ising models on the honeycomb, diced and decorated honeycomb lattices [3], as well as on the decorated square, triangular and diced lattices [4] were exactly treated long ago. Notice that later on these mapping transformations were further generalized in order to account for the single-ion anisotropy effect. The effect of uniaxial single-ion anisotropy on magnetic properties of the mixed-spin systems has been precisely investigated on the honeycomb lattice [5] and on some decorated planar lattices [6]. Moreover, a more general form of the star-triangle transformation regarding the biaxial single-ion anisotropy has recently been employed to obtain exact results for the mixed-spin Ising model on the honeycomb lattice [7]. To the best of our knowledge, these are the only mixed-spin systems with generally known exact solutions except the ones on the Bethe (Cayley tree) lattices, that can be studied within a discrete non-linear map [8] and/or using an approach based on exact recursion equations [9]. Among the remarkable models for which a precise solution is restricted to a certain subspace of interaction parameters only, one should further mention the mixed-spin Ising models on the union jack (centered square) lattice that can be mapped onto the eight-vertex model [10].

The purpose of this work is to provide an exact formulation of the mixed spin-1/2 and spin-1 Ising model on the diced lattice and to establish accurate phase diagrams of this system depending on the strength of the uniaxial and biaxial single-ion anisotropies as well as on the strength of the next-nearest-neighbour interaction. Exact results of this system are obtained by applying the generalized star-triangle transformation constituting an exact correspondence with the effective spin-1/2 Ising model on the triangular lattice. Finally, the effect of single-ion anisotropy terms and the next-nearest-neighbour interaction on the temperature dependence of the resultant magnetization will be clarified for the case of ferrimagnetic system.

The organization of this paper is as follows. In section 2, a detailed formula-

tion of the model system is presented and subsequently, exact expressions for the magnetization and critical temperatures are derived by establishing an exact correspondence with the effective spin-1/2 Ising model on the triangular lattice. The most interesting numerical results are presented and particularly discussed in section 3. Finally, some concluding remarks are given in section 4.

2. Model and its exact solution

Let us start with considering the mixed spin-1/2 and spin-1 Ising model on the diced lattice schematically illustrated on the left-hand side of figure 1. The mixed-spin diced lattice consists of two interpenetrating sub-lattices A and B which are formed by the spin-1/2 (empty circles) and spin-1 (solid circles) atoms, respectively. The Ising Hamiltonian defined upon the underlying lattice reads:

$$\hat{\mathcal{H}}_{\text{diced}} = -J \sum_{(i,j) \subset \mathcal{L}} \hat{S}_i^z \hat{\sigma}_j^z - J' \sum_{(k,l) \subset \mathcal{K}} \hat{\sigma}_k^z \hat{\sigma}_l^z - D \sum_{i=1}^{2N} (\hat{S}_i^z)^2 - E \sum_{i=1}^{2N} [(\hat{S}_i^x)^2 - (\hat{S}_i^y)^2], \quad (1)$$

where $\hat{\sigma}_j^z$ and \hat{S}_i^α ($\alpha = x, y, z$) stand for the spatial components of spin-1/2 and spin-1 operators, respectively, J denotes the exchange interaction between the nearest-neighbouring spin pairs and J' denotes the interaction between the next-nearest-neighbouring spin-1/2 atoms. Finally, the parameters D and E measure a strength of the uniaxial and biaxial single-ion anisotropies acting on the spin-1 sites and N denotes the total number of the spin-1/2 sites.




Figure 1. Diagrammatic representation of the diced lattice consisting of the mixed spin-1/2 (empty circles) and spin-1 (solid circles) sites, respectively. The solid (broken) lines depict the nearest-neighbour (next-nearest-neighbour) interactions. Using a star-triangle transformation, the mixed-spin diced lattice is mapped onto a simple spin-1/2 triangular lattice.

It can be clearly seen from figure 1 that the considered diced lattice belongs to loose-packed lattices, i.e. it can be decomposed into two interpenetrating sub-lattices A and B in such a way that all nearest neighbours of the spin-1 sites are the spin-1/2 sites and vice versa. Consequently, a trace over spin degrees of freedom of all spin-1 atoms can be performed before summing over all available configurations of the spin-1/2 atoms. It is therefore very convenient to write the total Hamiltonian

(1) as a sum of the commuting site Hamiltonians $\hat{\mathcal{H}}_{\text{diced}} = \sum_i \hat{\mathcal{H}}_i$, where each site Hamiltonian $\hat{\mathcal{H}}_i$ involves all interaction terms associated with one appropriate spin-1 site residing i^{th} lattice point of the sub-lattice B :

$$\begin{aligned} \hat{\mathcal{H}}_i = & - \hat{S}_i^z J (\hat{\sigma}_{i1}^z + \hat{\sigma}_{i2}^z + \hat{\sigma}_{i3}^z) - (\hat{S}_i^z)^2 D - [(\hat{S}_i^x)^2 - (\hat{S}_i^y)^2] E \\ & - J' (\hat{\sigma}_{i1}^z \hat{\sigma}_{i2}^z + \hat{\sigma}_{i2}^z \hat{\sigma}_{i3}^z + \hat{\sigma}_{i3}^z \hat{\sigma}_{i1}^z) / 2. \end{aligned} \quad (2)$$

The factor $1/2$ emerging in the last expression arises in order to avoid a double counting of each next-nearest-neighbour interaction when summing over all spin-1 sites. On account of the commutability between the site Hamiltonians (2), the partition function of the mixed-spin diced lattice can be partially factorized to:

$$\mathcal{Z}_{\text{diced}} = \text{Tr}_{\{\sigma\}} \prod_{i=1}^{2N} \text{Tr}_{S_i} \exp(-\beta \hat{\mathcal{H}}_i), \quad (3)$$

where $\beta = 1/(k_B T)$, k_B is Boltzmann's constant, T stands for the absolute temperature, the symbol Tr_{S_i} means a trace over the spin degrees of freedom of i^{th} spin-1 atom of the sub-lattice B and the symbol $\text{Tr}_{\{\sigma\}}$ stands for a trace over all possible spin configurations of the spin-1/2 atoms on the sub-lattice A .

After performing elementary calculations (see for instance [7]), one readily finds an analytical expression for $\text{Tr}_{S_i} \exp(-\beta \hat{\mathcal{H}}_i)$. Moreover, an explicit form of the relevant trace immediately implies a possibility of performing the familiar star-triangle mapping transformation [3]:

$$\begin{aligned} \text{Tr}_{S_i} \exp(-\beta \hat{\mathcal{H}}_i) &= \exp[\beta J' (\sigma_{i1}^z \sigma_{i2}^z + \sigma_{i2}^z \sigma_{i3}^z + \sigma_{i3}^z \sigma_{i1}^z) / 2] \\ &\quad \times \left\{ 1 + 2 \exp(\beta D) \cosh \left[\beta \sqrt{J^2 (\sigma_{i1}^z + \sigma_{i2}^z + \sigma_{i3}^z)^2 + E^2} \right] \right\} \\ &= C \exp[\beta R (\sigma_{i1}^z \sigma_{i2}^z + \sigma_{i2}^z \sigma_{i3}^z + \sigma_{i3}^z \sigma_{i1}^z) / 2] \end{aligned} \quad (4)$$

which establishes an exact correspondence between the mixed-spin Ising model on the diced lattice and its equivalent spin-1/2 Ising model on the triangular lattice provided that the transformation (4) is applied to all spin-1 sites (see figure 1). Notice that the unknown mapping parameters C and R are unambiguously given directly by the mapping transformation (4), which should hold for all available configurations of $\sigma_{i1}^z, \sigma_{i2}^z, \sigma_{i3}^z$ spins. According to this, one easily finds:

$$C^4 = V_1 V_2^3, \quad \beta R = \beta J' + 2 \ln(V_1/V_2), \quad (5)$$

with the expressions V_1 and V_2 defined as:

$$V_1 = 1 + 2 \exp(\beta D) \cosh \left(\beta \sqrt{(3J/2)^2 + E^2} \right), \quad (6)$$

$$V_2 = 1 + 2 \exp(\beta D) \cosh \left(\beta \sqrt{(J/2)^2 + E^2} \right). \quad (7)$$

Now, let us substitute the mapping relation (4) into the formula (3) gained for the partition function of the mixed-spin diced lattice. After straightforward rearrangement, one easily obtains an exact relationship between the partition function

$\mathcal{Z}_{\text{diced}}$ of the mixed-spin Ising model on the diced lattice and respectively, the partition function $\mathcal{Z}_{\text{triang}}$ of its corresponding spin-1/2 Ising model on the triangular lattice:

$$\mathcal{Z}_{\text{diced}}(\beta, J, J', D, E) = C^{2N} \mathcal{Z}_{\text{triang}}(\beta, R). \quad (8)$$

With regard to this, the mixed-spin diced lattice is mapped onto the spin-1/2 Ising model on the triangular lattice exactly solved several years ago [11–13].

The mapping relation (8) between the partition functions $\mathcal{Z}_{\text{diced}}$ and $\mathcal{Z}_{\text{triang}}$ represents a central result of our calculation. In fact, this relation can be utilized for establishing similar mapping relations for other important quantities such as Gibbs free energy, internal energy, magnetization, correlation functions, specific heat, etc. When combining the formula (8) with commonly used mapping theorems [14], one readily proves a validity of the following relations:

$$m_A \equiv \langle \hat{\sigma}_i^z \rangle_{\text{diced}} = \langle \hat{\sigma}_i^z \rangle_{\text{triang}} \equiv m_{\text{triang}}, \quad (9)$$

$$t_A \equiv \langle \hat{\sigma}_i^z \hat{\sigma}_j^z \hat{\sigma}_k^z \rangle_{\text{diced}} = \langle \hat{\sigma}_i^z \hat{\sigma}_j^z \hat{\sigma}_k^z \rangle_{\text{triang}} \equiv t_{\text{triang}}, \quad (10)$$

$$m_B \equiv \langle S_i^z \rangle_{\text{diced}} = 3m_A[F(3) + F(1)]/2 + 2t_A[F(3) - 3F(1)], \quad (11)$$

where the symbols $\langle \cdots \rangle_{\text{diced}}$ and $\langle \cdots \rangle_{\text{triang}}$ denote standard canonical average performed over the Ising models on the mixed spin-1/2 and spin-1 diced lattice and on the spin-1/2 triangular lattice, respectively, and the function $F(x)$ stands for:

$$F(x) = \frac{Jx/2}{\sqrt{(Jx/2)^2 + E^2}} \frac{2 \sinh(\beta \sqrt{(Jx/2)^2 + E^2})}{\exp(-\beta D) + 2 \cosh(\beta \sqrt{(Jx/2)^2 + E^2})}. \quad (12)$$

It should be pointed out that the set of equations (9)–(11) makes possible a straightforward calculation of both sub-lattice magnetization m_A and m_B of the mixed-spin system on the diced lattice. According to equation (9), the sub-lattice magnetization m_A directly equals to the corresponding magnetization m_{triang} of the triangular lattice, which is unambiguously given by the mapping relation (5). Note that the exact result for magnetization of the spin-1/2 Ising model on the triangular lattice was originally derived by Potts in his short note [12]:

$$m_{\text{triang}} = \frac{1}{2} \left[1 - \frac{16x^3}{(1+3x)(1-x)^3} \right]^{1/8}, \quad (13)$$

where $x = \exp(-\beta R)$. On the other hand, an exact knowledge of the triplet order parameter t_{triang} is required in order to compute the sub-lattice magnetization m_B in addition to the single-site magnetization m_{triang} . Exact results of Baxter and Choy [13] for the anisotropic triangular Ising model include, as specialization, the following exact solution for the triplet correlation of the isotropic triangular lattice:

$$t_{\text{triang}} = \frac{m_{\text{triang}}}{4} \left[1 + 2 \frac{x^2 - 2x^{-2} + 1 - \sqrt{(x^2 + 3)(x - 1)}}{(x - x^{-1})^2} \right]. \quad (14)$$

In view of the results (13) and (14), an exact solution of both sub-lattice magnetizations of the mixed-spin system is thus formally completed.

Finally, let us derive an analytical condition that determines a criticality of the mixed-spin diced lattice. It can be easily understood that both sub-lattice magnetizations m_A and m_B vanish if the magnetization m_{triang} of the corresponding triangular lattice tends to zero. It is therefore sufficient to substitute an exact critical point of the isotropic triangular lattice [11] into the mapping relation (5) in order to obtain an accurate condition for the criticality of the mixed-spin Ising model:

$$\beta_c J' + 2 \ln \left\{ \frac{1 + 2 \exp(\beta_c D) \cosh\left(\beta_c \sqrt{(3J/2)^2 + E^2}\right)}{1 + 2 \exp(\beta_c D) \cosh\left(\beta_c \sqrt{(J/2)^2 + E^2}\right)} \right\} = \ln 3, \quad (15)$$

where $\beta_c = 1/(k_B T_c)$ and T_c denotes the critical temperature.

3. Numerical results and discussion

Before proceeding to a discussion of the most interesting numerical results it is worthwhile mentioning that the results obtained in the preceding section hold for the ferromagnetic ($J > 0$) as well as antiferromagnetic ($J < 0$) nearest-neighbour interaction J . For simplicity, we shall restrict ourselves in what follows to the latter case corresponding to the ferrimagnetically ordered system. It is noteworthy, however, that the critical frontiers of the ferromagnetic model are congruent with those displayed below for the ferrimagnetic model as a result of an invariance of the critical condition (15) with respect to the transformation $J \rightarrow -J$. It should be also mentioned that the critical condition (15) does not depend on the sign of the parameter E . Hence, one may also consider $E \geq 0$ without loss of generality.

3.1. Ground-state phase diagrams

Let us first focus on the ground-state phase diagrams of the system under investigation. As the easy-plane uniaxial single-ion anisotropy ($D < 0$) reinforces, a first-order phase transition from the ferrimagnetically ordered phase towards the disordered phase should be expected to occur owing to energetic favoring of a non-magnetic state $S = 0$ of the spin-1 atoms. It is quite interesting to ascertain, however, that the biaxial single-ion anisotropy stabilizes the ferrimagnetically ordered ground state as far as it enhances the critical value of the uniaxial single-ion anisotropy at which both ferrimagnetic and disordered phases coexist together:

$$\frac{D_c}{|J|} = \frac{-3}{2\sqrt{1 - (E/D)^2}}. \quad (16)$$

This remarkable feature turns out to be in an obvious contradiction with the naive expectations, since the biaxial term E entails uprise of local quantum fluctuations that manifest themselves in a quantum reduction of the sub-lattice magnetization

m_B . Actually, the arbitrary but non-zero biaxial single-ion anisotropy destroys a perfect ferrimagnetic order even at zero temperature and the present quantum ferrimagnetic phase can be characterized through its sub-lattice order parameters given by:

$$m_A = -\frac{1}{2}, \quad m_B = \frac{3}{\sqrt{9 + 4(E/|J|)^2}}. \quad (17)$$

As far as the effect of next-nearest-neighbour interaction J' is concerned, one should vigorously distinguish between the effect of ferromagnetic ($J' > 0$) and antiferromagnetic ($J' < 0$) interaction. The former interaction aligns ferromagnetically the next-nearest-neighbouring spin-1/2 atoms and thus, the spin system remains long-range-ordered independently of the strength of the uniaxial single-ion anisotropy that acts purely on the spin-1 sites. On the other hand, the latter interaction causes a spin frustration between the next-nearest-neighbours and consequently it behaves in compliance with the uniaxial single-ion anisotropy in view of destroying the ferrimagnetic long-range order. This synergetic effect can also be viewed from a modified version of the critical condition (16) when switching on the next-nearest-neighbour interaction J' :

$$\frac{D_c}{|J|} = \frac{-3}{2\sqrt{1 - (E/D)^2}} - \frac{J'}{2|J|} \quad (\text{for } J' < 0 \text{ only}). \quad (18)$$

3.2. Finite-temperature phase diagrams

Now, let us turn to a detailed discussion dealing with the finite temperature phase diagrams. For this purpose, figure 2 shows the variation of reduced critical temperature with the uniaxial single-ion anisotropy for several fixed values of the ratio E/D and $J'/|J| = 0.0$.




Figure 2. The reduced critical temperature against the uniaxial single-ion anisotropy at $E/D = 0.0, 0.2, 0.3, 0.4, 0.5$ in ascending order along the direction of arrows. The case $E/D = 0.0$ is plotted as a dashed line for visual effect.

The quantum ferrimagnetic phase can be located inside the depicted phase boundaries, while above them the usual paramagnetic phase becomes stable. There are two interesting things to observe here. Firstly, there is a slight suppression of the critical temperature in the region of easy-axis uniaxial single-ion anisotropies ($D > 0$) for any $E \neq 0$. This unusual feature can be interpreted in terms of strengthening the quantum fluctuations caused by the non-zero biaxial single-ion anisotropy. Actually, the greater the ratio E/D , the more impressive reduction of T_c can be observed. Secondly, the insert of figure 2 clearly demonstrates that the system exhibits

interesting reentrance in the vicinity of the critical value $D_c/|J|$ given by the condition (16). For the uniaxial anisotropies slightly below this critical value, one starts from the disordered ground state before entering the ordered phase at lower critical temperature and finally, the ferrimagnetic long-range order disappears due to thermal fluctuations at the upper critical temperature. Notice that such pure reentrant transition with two consecutive critical points have not been reported in the analogous mixed-spin system on the honeycomb lattice [7]. This observation would suggest that a presence of the reentrant transition can possibly be attributed to higher coordination number of the spin-1/2 sites.

The influence of the next-nearest-neighbour interaction on the phase diagrams is illustrated in figure 3. As one can see, the spin system remains long-range ordered for arbitrary strength of the ferromagnetic next-nearest-neighbour interaction ($J' > 0$). When small ferromagnetic coupling J' is taken into consideration, however, the critical temperature versus uniaxial anisotropy dependence can be characterized by the *S*-shaped curve indicating an occurrence of the reentrant transitions with three consecutive critical points (see the insert in figure 3). It is noteworthy that this reentrant behaviour gradually vanishes as the ratio $J'/|J|$ strengthens. Even more interesting situation emerges when assuming the antiferromagnetic next-nearest-neighbour interaction ($J' < 0$). In the latter case, the displayed critical lines imply an existence of reentrant transitions with either two or three consecutive critical points before approaching $D_c/|J|$ given by equation (18). Notice that a range of the critical boundaries corresponding to the reentrant transition with two (three) critical temperatures overwhelms at relatively smaller (greater) values of $|J'|/|J|$, while the reentrance completely disappears from the critical lines above certain boundary value of $|J'|/|J|$. Finally, it is worth mentioning that similar influence of the next-nearest-neighbour interaction on the criticality is observable even if a non-zero strength of the biaxial single-ion anisotropy would be considered.




Figure 3. The reduced critical temperature against the uniaxial single-ion anisotropy at $J'/|J| = -0.2, -0.1, -0.05, -0.025, 0.0, 0.025, 0.05, 0.1, 0.2$ in ascending order along the direction of arrows. The case $J'/|J| = 0.0$ is plotted as a dashed line for visual effect. A dotted line shows the line of compensation temperatures.

3.3. Temperature dependence of magnetization

In this part, the temperature dependence of magnetization (to be further abbreviated as $m(T)$ curves) will be explored in particular. When both kinds of atoms are stoichiometrically taken into account, the resultant magnetization reduced per

one lattice site of the mixed-spin diced lattice can be defined as $m = |m_A + 2m_B|/3$. Figure 4(a) depicts the effect of uniaxial single-ion anisotropy on the shape of $m(T)$ curves. The downward curvature observed for strong negative uniaxial anisotropies




Figure 4. Temperature variation of the resultant magnetization as the uniaxial single-ion anisotropy varies and the ratio $J'/|J| = 0.0$. The parameter of biaxial single-ion anisotropy is fixed to be: (a) $E/D = 0.0$; (b) $E/D = 0.2$.




Figure 5. The effect of small ferromagnetic next-nearest-neighbour interaction $J'/|J| = 0.05$ on the temperature dependence of magnetization. Figure 5(a) shows the variation of the resultant magnetization, figure 5(b) depicts in addition to the total magnetization (solid lines), the temperature variation of the sub-lattice magnetization m_A (dotted lines) and m_B (dashed lines), respectively.

can obviously be attributed to a more rapid thermal variation of the sub-lattice magnetization m_B , which is thermally more easily disturbed than m_A under the condition $D < 0$. In accordance with the phase diagrams displayed earlier, one finds $m(T)$ curves exhibiting the reentrant transitions with two consecutive critical points slightly below the critical value $D_c/|J|$ given by the condition (16). For comparison, figure 4(b) illustrates how the biaxial single-ion anisotropy affects a temperature variation of the resultant magnetization when the ratio $E/D = 0.2$ is fixed and $D/|J|$ varies. It is quite evident that the most ultimate difference between

the $m(T)$ curves depicted in figures 4(a) and 4(b), respectively, rests in the zero-temperature limit of the latter ones that do not start from their saturation value owing to the quantum reduction of the sub-lattice magnetization m_B caused by the biaxial anisotropy.

Next, the effect of next-nearest-neighbour interaction J' upon the shape of $m(T)$ curves will be clarified. The effect of biaxial anisotropy is neglected here for simplicity, however, it worthy to note that the curves displayed below remain qualitatively unchanged even if the biaxial anisotropy term would be non-zero. Figure 5(a) shows typical $m(T)$ curves under the assumption of small ferromagnetic interaction $J'/|J| = 0.05$. The most remarkable differences compared to the $m(T)$ curves discussed earlier can be found in the vicinity of the boundary value $D_c/|J| = -1.5$, which is strong enough to force all the spin-1 atoms towards their non-magnetic state $S = 0$ in the ground state. For illustration, figure 5(b) depicts the sub-lattice magnetization of two most striking dependencies in addition to the resultant magnetization. As one can see, there appears $m(T)$ curve with one compensation point if the uniaxial anisotropy is chosen slightly below its boundary value (for instance $D/|J| = -1.6$). The compensation phenomenon arises on account of thermally induced increase of the sub-lattice magnetization m_B , which overwhelms the sub-lattice magnetization m_A above the compensation temperature. When the uniaxial anisotropy is taken to be even more negative, there emerges $m(T)$ curve with three consecutive critical points instead of the one with the compensation point (the case $D/|J| = -1.63$). Notice that a further decrease of the uniaxial single-ion anisotropy shrinks the temperature range between the second and third critical point until the ordered phase completely vanishes from this higher-temperature region.




Figure 6. The same as in figure 5 but for the case of small antiferromagnetic next-nearest-neighbour interaction $J'/|J| = -0.05$.

At last, we shall briefly discuss the effect of small antiferromagnetic next-nearest-neighbour interaction ($J'/|J| = -0.05$) on the temperature dependence of the magnetization as shown in figure 6. According to equation (18), the critical line ter-

minates due to the spin frustration already at $D_c/|J| = -1.475$. It is quite apparent from figures 6(a) and 6(b) that the magnetization exhibits two reentrant transitions for uniaxial anisotropies slightly below this critical value, for example at $D/|J| = -1.5$. Contrary to this, the resultant magnetization may show the reentrant transition with three critical temperatures when the uniaxial anisotropy is selected slightly above this boundary value (the case $D/|J| = -1.47$). Whether the reentrant transition with two or three critical points is observed, however, depends very sensitively on the relative magnitude of all interaction parameters. It appears worthwhile remarking that both kinds of reentrant transitions disappear as the magnitude of next-nearest-neighbour interaction is considered to be strong enough.

4. Concluding remarks

In the present paper, the mixed spin-1/2 and spin-1 Ising model on the diced lattice has been exactly treated within the generalized star-triangle mapping transformation. The main focus of the present work is aimed at the effect of the next-nearest-neighbour interaction, uniaxial and biaxial single-ion anisotropies on the magnetic behaviour of the system under investigation. It has been found that the single-ion anisotropy may possibly cause the reentrant transition with two critical temperatures only. On the other hand, the ferromagnetic next-nearest-neighbour interaction may lead just to the reentrant transition with three consecutive critical points, whereas the antiferromagnetic next-nearest-neighbour coupling may be considered as a possible cause of both kinds of reentrant transitions.

Another interesting finding to emerge here consists in the fact that the biaxial single-ion anisotropy is confirmed to be a possible source of the quantum reduction of magnetization. Our further observations suggest that the biaxial anisotropy cannot qualitatively change the character of the reentrant transition caused by the uniaxial single-ion anisotropy term or the next-nearest-neighbour coupling. Finally, we should remark that a combined effect of uniaxial and biaxial single-ion anisotropies with the next-nearest-neighbour interaction is responsible for manifold temperature dependencies of the resultant magnetization by assuming the ferrimagnetically ordered spin system.

References

1. Onsager L., Phys. Rev., 1944, **65**, 117.
2. Baxter R. J. Exactly Solved Models in Statistical Mechanics. Academic Press, New York, 1982.
3. Fisher M.E., Phys. Rev., 1959, **113**, 969; Szyzi I. Phase Transitions and Critical Phenomena, vol. 1, eds. Domb C. and Green M.S. Academic, New York, 1972.
4. Yamada K., Progr. Theor. Phys., 1969, **42**, 1106.
5. Gonçalves L.L., Phys. Scripta, 1985, **32**, 248; Gonçalves L.L., Phys. Scripta, 1986, **33**, 192; Gonçalves L.L., Phys. Scripta, 1987, **35**, 105; Tucker J.W., J. Magn. Magn.

- Mater., 1999, **95**, 133; Dakhama A., Benayad N., J. Magn. Magn. Mater., 2000, **231**, 117.
6. Jaščur M., Physica A, 1998, **252**, 217; Dakhama A., Physica A, 1998, **252**, 225; Lacková S., Jaščur M., Acta Phys. Slovaca, 1998, **48**, 623.
 7. Strečka J., Jaščur M., Acta Electrotechnica et Informatica, 2002, **2**, 102; Strečka J., Jaščur M., Phys. Rev. B, 2004, **70**, 014404; Jaščur M., Strečka J., Physica A, in press.
 8. da Silva N. R., Salinas S.R., Phys. Rev. B, 1991, **44**, 852.
 9. Albayrak E., Keskin M., J. Magn. Magn. Mater., 2003, **261**, 196; Albayrak E., Int. J. Mod. Phys. B, 2004, **18**, 3959; Albayrak E., Alci A., Physica A, 2005, **345**, 48.
 10. Lipowski A., Horiguchi T., J. Phys. A: Math. Gen., 1995, **28**, L261; Strečka J., to be submitted to Condens. Matter Phys., 2005.
 11. Houtappel R.M.F., Physica (Amsterdam), 1950, **16**, 425; Temperley H.N.V., Proc. R. Soc. London, Ser. A, 1950, **203**, 202; Wannier G.H., Phys. Rev., 1950, **79**, 357; Newell G.F., Phys. Rev., 1950, **79**, 876.
 12. Potts R.B., Phys. Rev., **88**, 352.
 13. Baxter R.J., Choy T.C., Proc. R. Soc. London, Ser. A, 1989, **423**, 279.
 14. Barry J.H., Khatun M., Tanaka T., Phys. Rev. B, 1988, **37**, 5193; Khatun M., Barry J.H., Tanaka T., Phys. Rev. B 1990, **42**, 4398; Barry J.H., Tanaka T., Khatun M., Múnera C.H., Phys. Rev. B, 1991, **44**, 2595; Barry J.H., Khatun M., Phys. Rev. B, 1995, **51**, 5840.

Переходи повернення для моделі Ізінга зі змішаними спінами на гратці типу грального кубіка

М.Ящур, Й.Стречка

Відділ теоретичної фізики та астрофізики,
факультет науки, Університет ім. П.Й.Шафаріка,
040 01 Кошіце, Словацька Республіка

Отримано 18 липня, 2005

Магнітна поведінка змішаної спін-1/2 та спін-1 моделі Ізінга на гратці типу грального кубіка досліджується з використанням точного перетворення співставлення зірка-трикутник. Отримано, що як уніаксіальна так і біаксіальна анізотропія іона, діючи на вузли зі спіном-1, потенційно можуть спричинити перехід повернення з двома послідовними критичними точками. На протипагу цьому, ефект взаємодії наступних за найближчими сусідами між вузлами зі спінами-1/2 ймовірно приводить до спадного переходу з трьома критичними температурами додатково до переходу з лише двома критичними точками. Форма залежності повного намагнічення від температури досліджується, зокрема, для випадку ферімагнітно впорядкованих систем.

Ключові слова: *перехід повернення, модель Ізінга, точний розв'язок, перетворення зірка-трикутник, уніаксіальна та біаксіальна іонна анізотропія*

PACS: 75.10.Nk, 05.50.+q, 75.40