

EVALUATION OF SERUM LEVELS OF MULTIPLE CYTOKINES AND ADHESION MOLECULES IN PATIENTS WITH NEWLY DIAGNOSED ACUTE LYMPHOBLASTIC LEUKEMIA USING BIOCHIP ARRAY TECHNOLOGY

J.M. Horacek^{1,2,*}, T. Kupsa^{1,2}, M. Vasatova³, L. Jebavy^{1,2}, P. Zak²

¹Department of Internal Medicine, University of Defence, Faculty of Military Health Sciences, Hradec Kralove, Czech Republic

²4th Department of Internal Medicine — Hematology, University Hospital and Charles University, Faculty of Medicine, Hradec Kralove, Czech Republic

³Institute of Clinical Biochemistry and Diagnostics, University Hospital, Hradec Kralove, Czech Republic

Aim: Evaluation of serum levels of 17 cytokines and 5 adhesion molecules in patients with acute lymphoblastic leukemia (ALL) and in healthy subjects using biochip array technology. This approach allows multi-analytical determination from a single sample.

Methods: A total of 15 ALL patients and 15 healthy subjects (blood donors) were studied. Serum samples were analyzed by biochip based immunoassays on the Evidence Investigator analyzer. T-tests were used for statistical analysis. **Results:** Comparing cytokine and adhesion molecule levels in ALL patients and in healthy subject, we found significant increase in serum VCAM-1 ($p < 0.000001$), ICAM-1 ($p < 0.0001$), L-selectin ($p < 0.0001$), IL-8 ($p < 0.001$), MCP-1 ($p < 0.01$), and significant decrease ($p < 0.01$) in serum IL-3 and IL-4. **Conclusion:** Our results indicate that serum levels of specific cytokines and adhesion molecules (VCAM-1, ICAM-1, L-selectin, IL-8, IL-3, IL-4, MCP-1) are significantly altered in patients with newly diagnosed ALL, reflecting activity of the disease. Further investigation is needed to establish if these alterations could be used as a prognostic indicator for ALL.

Key Words: cytokines, adhesion molecules, biochip array, acute lymphoblastic leukemia.

Cytokines and adhesion molecules have been studied in many pathological states including hematological malignancies [1–3] and acute leukemias, both myeloid (AML) and lymphoblastic (ALL) [4, 5]. Alterations in this interacting functional network may have direct effect on the malignant cells or have indirect effect on leukemogenesis through altered functions of bone marrow stromal elements [6, 7]. The knowledge gained from multi-analytical determination of cytokines and adhesion molecules could allow better diagnosis and management of hematological malignancies, since cytokines or their receptors may also represent a target for specific anticancer therapy at the molecular level. Recently, some studies reported the possible diagnostic and prognostic use of cytokine levels in newly diagnosed acute leukemias and myelodysplastic syndromes [8–11].

The aim of our pilot study was to evaluate serum levels of multiple cytokines and adhesion molecules in patients with newly diagnosed ALL and in healthy subjects using the innovative biochip array technology. This generates a patient profile, which is relevant when investigating interacting functional networks.

Received: August 16, 2013.

*Correspondence: E-mail: horacek@pmfhk.cz

Abbreviations used: ALL – acute lymphoblastic leukemia; AML – acute myeloid leukemia; CR – complete remission; EGF – epidermal growth factor; ICAM-1 – intercellular adhesion molecule-1; IFN-gamma – interferon-gamma; IL – interleukin; MCP-1 – monocyte chemotactic protein-1; TNF-alpha – tumor necrosis factor-alpha; VCAM-1 – vascular cell adhesion molecule-1; VEGF – vascular endothelial growth factor.

Subjects. A total of 15 newly diagnosed ALL patients (median age 46, range 24–63 years, 11 males) and 15 healthy subjects (median age 41, range 25–58 years, 11 males) were studied. The study was approved by the local Ethics Committee and all patients gave a written consent.

Multi-analytical evaluation. We evaluated circulating levels of the following 17 cytokines and 5 adhesion molecules: interleukins (IL-1 alpha, IL-1 beta, IL-2, IL-3, IL-4, IL-6, IL-7, IL-8, IL-10, IL-12p70, IL-13, IL-23), vascular endothelial growth factor (VEGF), tumor necrosis factor-alpha (TNF-alpha), interferon-gamma (IFN-gamma), epidermal growth factor (EGF), monocyte chemotactic protein-1 (MCP-1), E-selectin, L-selectin, P-selectin, intercellular adhesion molecule-1 (ICAM-1), vascular cell adhesion molecule-1 (VCAM-1). All analytes were measured by biochip array technology using chemiluminescent sandwich immunoassays applied to the Evidence Investigator analyzer (“Randox Laboratories Ltd.”, Crumlin, UK). We analyzed serum samples at the diagnosis of ALL (active leukemia) and in healthy subjects (blood donors).

Statistical analysis. Statistical analysis was performed with the “Statistica” program. T-tests were used. The values were expressed as mean \pm SD. Probability values (p) < 0.01 were considered statistically significant.

In newly diagnosed ALL patients, we found significant increase in serum VCAM-1 (1078.54 ± 456.96 mcg/L vs. 328.31 ± 88.66 mcg/L; $p < 0.000001$), ICAM-1 (499.57 ± 237.53 mcg/L vs. 196.69 ± 36.06 mcg/L; $p < 0.0001$), L-selectin (2366.33 ± 1035.37 mcg/L vs. $1104.54 \pm$

243.45mcg/L; $p < 0.0001$), IL-8 (34.07 ± 28.52 ng/L vs. 4.87 ± 3.09 ng/L; $p < 0.001$), MCP-1 (433.99 ± 328.59 ng/L vs. 153.25 ± 53.60 ng/L; $p < 0.01$). On the other hand, we found significant decrease in serum IL-3 (7.34 ± 3.41 ng/L vs. 11.53 ± 4.66 ng/L; $p < 0.01$), IL-4 (1.10 ± 1.08 ng/L vs. 3.27 ± 2.21 ng/L; $p < 0.01$). No significant differences were found in the levels of other evaluated cytokines and adhesion molecules.

To our knowledge, this is the first published study using the innovative biochip array technology to determine circulating levels of cytokines and adhesion molecules in ALL patients.

Altered levels of cytokines and adhesion molecules have been found in many pathological states and have been linked to many diseases such as autoimmune diseases, allergies, cardiovascular diseases and cancer [12–17]. The cytokine system constitutes an interacting functional network where the contribution from single cytokines is modulated by the levels of other cytokines. It may therefore be more relevant to look at the total serum profile of cytokines and adhesion molecules.

Biochip array technology enables simultaneous detection of multiple cytokines and adhesion molecules in a single sample and provides valuable information relating to each tested analyte and possible associations between analytes in each sample [18, 19].

Our results indicate that serum levels of specific cytokines and adhesion molecules (VCAM-1, ICAM-1, L-selectin, IL-8, IL-3, IL-4, MCP-1) are significantly altered in patients with newly diagnosed ALL, reflecting activity of the disease. Further investigation is needed to establish if the alterations observed in the levels of these molecules could be used as a prognostic indicator for ALL.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

The work was supported by a long-term organization development plan 1011 (Faculty of Military Health Sciences, Hradec Kralove) and by a specific research project "Analysis of defined prognostic factors in acute leukemia" (Faculty of Military Health Sciences, Hradec Kralove).

REFERENCES

1. Bruserud Ø, Kittang AO. The chemokine system in experimental and clinical hematology. *Curr Top Microbiol Immunol* 2010; **341**: 3–12.
2. Mellgren K, Hedegaard CJ, Schmiegelow K, *et al.* Plasma cytokine profiles at diagnosis in pediatric patients with non-hodgkin lymphoma. *J Pediatr Hematol Oncol* 2012; **34**: 271–5.
3. Deeg HJ. Cytokines in graft-versus-host disease and the graft-versus-leukemia reaction. *Int J Hematol* 2001; **74**: 26–32.

4. Lowenberg B, Touw IP. Hematopoietic growth factors and their receptors in acute leukemia. *Blood* 1993; **81**: 281–92.

5. Kupsa T, Horacek JM, Jebavy L. The role of cytokines in acute myeloid leukemia: a systematic review. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub* 2012; **156**: 291–301.

6. Konopleva MY, Jordan CT. Leukemia stem cells and microenvironment: biology and therapeutic targeting. *Clin Oncol* 2011; **29**: 591–9.

7. Reikvam H, Hatfield KJ, Fredly H, *et al.* The angioregulatory cytokine network in human acute myeloid leukemia — from leukemogenesis via remission induction to stem cell transplantation. *Eur Cytokine Netw* 2012; **23**: 140–53.

8. Tsimberidou AM, Estey E, Wen S, *et al.* The prognostic significance of cytokine levels in newly diagnosed acute myeloid leukemia and high-risk myelodysplastic syndromes. *Cancer* 2008; **113**: 1605–13.

9. Kornblau SM, McCue D, Singh N, *et al.* Recurrent expression signatures of cytokines and chemokines are present and are independently prognostic in acute myelogenous leukemia and myelodysplasia. *Blood* 2010; **116**: 4251–61.

10. Leblebisatan G, Antmen B, Saşmaz I, *et al.* Vascular endothelial growth factor levels in childhood acute lymphoblastic and myeloblastic leukemia. *Indian J Hematol Blood Transfus* 2012; **28**: 24–8.

11. Fung FY, Li M, Breunis H, *et al.* Correlation between cytokine levels and changes in fatigue and quality of life in patients with acute myeloid leukemia. *Leuk Res* 2013; **37**: 274–9.

12. Berrahmoune H, Lamont J, Fitzgerald P, *et al.* Inter-individual variation of inflammatory markers of cardiovascular risks and diseases. *Clin Chem Lab Med* 2005; **43**: 671–84.

13. Kavsak PA, Lee A, Hirte H, *et al.* Cytokine elevations in acute coronary syndrome and ovarian cancer: a mechanism for the up-regulation of the acute phase proteins in these different disease etiologies. *Clin Biochem* 2008; **41**: 607–10.

14. Poh YW, Gan SY, Tan EL. Effects of IL-6, IL-10 and TGF- β on the expression of survivin and apoptosis in nasopharyngeal carcinoma TW01 cells. *Exp Oncol* 2012; **34**: 85–9.

15. Coussens LM, Werb Z. Inflammation and cancer. *Nature* 2002; **420**: 860–7.

16. Dranoff G. Cytokines in cancer pathogenesis and cancer therapy. *Nat Rev Cancer* 2004; **4**: 11–22.

17. Naumnik W, Naumnik B, Niewiarowska K, *et al.* Novel cytokines: IL-27, IL-29, IL-31 and IL-33. Can they be useful in clinical practice at the time diagnosis of lung cancer? *Exp Oncol* 2012; **34**: 348–53.

18. McAleer D, McPhillips FM, FitzGerald SP, *et al.* Application of Evidence Investigator for the simultaneous measurement of soluble adhesion molecules: L-, P-, E-selectins, VCAM-1 and ICAM-1 in a biochip platform. *J Immunoassay Immunochem* 2006; **27**: 363–78.

19. Fitzgerald SP, McConnell RI, Huxley A. Simultaneous analysis of circulating human cytokines using a high-sensitivity cytokine biochip array. *J Proteome Res* 2008; **7**: 450–5.